

ANSWER TO A QUESTION OF ROSŁANOWSKI AND SHELAH

MÁRK POÓR

ABSTRACT. In [14] Rosłanowski and Shelah asked whether every locally compact non-discrete group has a null but non-meager subgroup, and conversely, whether it is consistent with *ZFC* that in every locally compact group a meager subgroup is always null. They gave affirmative answers for both questions in the case of the Cantor group and the reals. In this paper we give affirmative answers for the general case.

1. PRELIMINARIES AND NOTATIONS

If we state that topological groups G and H are isomorphic, in symbols $G \simeq H$, then we mean that there is an algebraic isomorphism, which is also a homeomorphism. Under the symbol $\leq$ we mean the subgroup relation, i.e. $H \leq G$ symbols that H is a subgroup of G , and if we write $H \triangleleft G$, then H is a normal subgroup of G . All topological groups are assumed to be Hausdorff. (It is known that for a locally compact Hausdorff topological group G , if $C \triangleleft G$ is a compact normal subgroup, then G/C is locally compact and Hausdorff.)

A topological group G is locally compact if for each $g \in G$, there is a neighborhood B of g , which is compact (i.e. each point has an open neighborhood which has compact closure). Since in compact Hausdorff spaces the Baire category theorem is true (i.e. no nonempty open set can be covered by countably many nowhere dense sets), it is also true in locally compact Hausdorff spaces.

For any topological space X , $\mathcal{B}(X)$ denotes the Borel sets, i.e. the σ -algebra generated by the open sets.

Recall the following definition of left Haar Borel measure:

Definition 1. Let G be a locally compact group, and μ be a Borel measure, i.e.

$$\mu : \mathcal{B}(G) \rightarrow [0, \infty]$$

such that

- μ is left-invariant, i.e. for every $g \in G$, $B \in \mathcal{B}(G)$

$$\mu(gB) = \mu(B),$$
- $\mu(U) > 0$ for each open $U \neq \emptyset$,
- $\mu(K) < \infty$ for each compact set K ,

The author was supported by the National Research, Development and Innovation Office – NKFIH, grant no. 104178.

- μ is inner regular with respect to the compact sets, that is for every B Borel

$$\mu(B) = \sup\{\mu(K) : K \subseteq B, K \text{ is compact}\}.$$

Then μ is called a left Haar Borel measure of G .

It is known that a left Haar Borel measure always exists, and it is unique up to a positive multiplicative constant.

Under left Haar measure we will mean the completion of a left Haar Borel measure.

Definition 2. Let G be a locally compact group. Suppose that ν is a left Haar Borel measure, and μ is the completion of ν , that is a subset $H \subseteq G$ is measurable if it differs from a Borel set by (at most) a null Borel set, i.e.

$$\text{if } (\exists B, B' \in \mathcal{B}(G) : B \triangle H \subseteq B' \wedge \nu(B') = 0) \text{ then } \mu(H) := \nu(B).$$

Then μ is a left Haar measure of G .

From now on for a locally compact group G μ_G will always symbol a left Haar measure.

It is easy to see (from the inner regularity) that if a set H is locally null, then it is null, i.e.

$$(\exists \text{ open cover } (U_\alpha)_{\alpha \in I} \text{ of } H : (\forall \alpha) \mu_G(H \cap U_\alpha) = 0) \Rightarrow \mu_G(H) = 0. \quad (1)$$

From now on $\mathcal{N}$ will denote the null-ideal, i.e.

$$\mathcal{N} = \{H \subseteq G : \mu(H) = 0\}.$$

Remark 3. There is another definition of Haar Borel measure, namely, instead of inner regularity wrt. the compact sets we can require

- outer regularity wrt. the open sets:

$$\mu(B) = \inf\{\mu(U) : B \subseteq U, U \text{ is open}\} \text{ for every } B \text{ Borel,}$$

- inner regularity wrt. the compact sets, only in case of open sets:

$$\mu(U) = \sup\{\mu(K) : K \subseteq U, K \text{ is compact}\} \text{ for every } U \text{ open}$$

which also exists and is unique up to a positive multiplicative constant. This definition results in a different left-invariant Borel measure only in the case of non- σ -compact groups. The proofs in the first chapter of this paper would also work with this definition of the Haar measure, but in the second we will make use of the property (1), and the theorem which we will present is simply not true if requiring these type of regularity properties, in Example 38 we mention a well-known counterexample.

Definition 4. Let $(I, \preceq)$ be a directed partially ordered set (poset), i.e. for every $a, b \in I$ there exists a common upper bound c . And let G_a ($a \in I$) be topological groups indexed by elements of I , and let $\varphi_{a,b} : G_b \rightarrow G_a$ for each pair $a \preceq b$ be surjective continuous homomorphisms, such that whenever $a \preceq b \preceq c$ then

$$\varphi_{a,c} = \varphi_{a,b} \circ \varphi_{b,c}.$$

Then this system is called an inverse system, and the inverse limit of the system $((G_a)_{a \in I}, (\varphi_{a,b})_{a \preceq b})$ is

$$\varprojlim_{i \in I} G_i = \left\{ (x_i)_{i \in I} \in \prod_{i \in I} G_i : x_a = \varphi_{a,b}(x_b) \ (\forall a \preceq b \in I) \right\}$$

with the topology inherited from the product space, and with the obvious (pointwise) group structure.

If $I = \{i\}$, then we identify $\varprojlim_{i \in I} G_i$ with G_i .

Remark 5. By the definition of the product and subspace topology, there is a sub-base consisting of sets of the form

$$\left(\varprojlim_{a \in I} G_a \right) \cap \left(U_a \times \prod_{b \in I \setminus \{a\}} G_b \right) \quad (a \in I, U_a \subseteq G_a \text{ open}).$$

But if $a \preceq d$ (since $(g_i)_{i \in I} \in \varprojlim_{a \in I} G_a \Rightarrow \varphi_{a,d}(x_d) = x_a$):

$$\left(\varprojlim_{i \in I} G_i \right) \cap \left(U_a \times \prod_{b \in I \setminus \{a\}} G_b \right) = \left(\varprojlim_{i \in I} G_i \right) \cap \left(\varphi_{a,d}^{-1}(U_a) \times \prod_{b \in I \setminus \{d\}} G_b \right).$$

Thus these sets are closed under finite intersection (assume that $a, c \preceq d$), indeed,

$$\begin{aligned} & \left(\left(\varprojlim_{i \in I} G_i \right) \cap \left(U_a \times \prod_{b \in I \setminus \{a\}} G_b \right) \right) \cap \left(\left(\varprojlim_{i \in I} G_i \right) \cap \left(U_c \times \prod_{b \in I \setminus \{c\}} G_b \right) \right) = \\ & \left(\left(\varprojlim_{i \in I} G_i \right) \cap \left(\varphi_{a,d}^{-1}(U_a) \times \prod_{b \in I \setminus \{d\}} G_b \right) \right) \cap \left(\left(\varprojlim_{i \in I} G_i \right) \cap \left(\varphi_{c,d}^{-1}(U_c) \times \prod_{b \in I \setminus \{d\}} G_b \right) \right) = \\ & = \left(\varprojlim_{i \in I} G_i \right) \cap \left(\left(\varphi_{c,d}^{-1}(U_c) \cap \varphi_{a,d}^{-1}(U_a) \right) \times \prod_{b \in I \setminus \{d\}} G_b \right). \end{aligned}$$

Hence, in fact, this sub-base is a base.

It is known that for any locally compact group the null-ideals of a left Haar measure and a right Haar measure coincide, thus we can speak about null sets ([8, 442F]).

We are interested in null, but non-meager, and conversely meager but non-null subgroups of locally compact groups. Only the non-discrete case is of interest, since non-empty open sets are of positive measure, thus in a discrete group $\{e\}$ has positive measure, and obviously is of second category. It is known that under the Continuum Hypothesis one can construct null but non-meager and meager but non-null subgroups in arbitrary non-discrete locally compact groups. In [14], Rosłanowski, and Shelah constructed *ZFC* examples for null, but non-meager subgroups of the Cantor group and the reals, and showed that it is consistent with *ZFC* that in the two groups every meager subgroup is null. Then they asked two questions

Problem 5.1 [14]

- (1) Does every locally compact group (with complete Haar measure) admit a null non-meager subgroup?
- (2) Is it consistent that no locally compact group has a meager non-null subgroup?

The fact, that the nonexistence of null, non-meager subgroups in the Cantor group is consistent was known earlier: Friedman proved that it is consistent with *ZFC* that every F_σ set in $2^\omega \times 2^\omega$ containing a rectangle of positive outer measure must contain a rectangle of positive measure. Shelah and Fremlin proved that this statement implies the nonexistence of meager, non-null subgroups in 2^ω (see [5]).

In the first part of this paper we will construct a null but non-meager subgroup in the case of the inverse limit of a countable sequence of finite groups, and in arbitrary second countable Lie-groups, and we show that these cases are sufficiently general.

In the second part, after noticing that in the model where Friedman's theorem holds, it holds not only for 2^ω but for every locally compact Polish group, by lemmas from the first part we reduce the case of arbitrary locally compact groups to the case of the Polish ones.

2. NULL, BUT NON-MEAGER SUBGROUPS

Using ideas from [14], we will construct a null but non-meager subgroup in inverse limits of countable many finite groups, and in second countable Lie-groups. First, from Lemma 7 to Lemma 22, we will state lemmas so that we can restrict ourselves to those sufficiently general cases. We will find an appropriate Lie group or profinite quotient for arbitrary locally compact groups, and we will show that the pull-back of a null but non-meager subgroup in that quotient is null and non-meager in the original group.

Remark 6. *Using ideas from [7], in the case of locally compact Polish groups we could much more easily reduce the general case to the case of Lie groups and profinite groups. Moreover, if considering only locally compact Abelian Polish groups, then we only would have to construct null but non-meager subgroups in $\mathbb{R}$, countable product of finite groups and the p -adic integers.*

After that, by Proposition 25 the profinite case is handled, and later there will be a similar construction for the Lie groups (Proposition 27). Finally we summarize, and state our main result, Theorem 29.

Recall the following lemma [15, Thm. 1.6.1].

Lemma 7. *(Gleason-Yamabe theorem, stronger version). Let G be a locally compact group. Then there exists an open subgroup G' of G such that, for any open neighborhood U of the identity in G' , there exists a compact normal subgroup $L_U \subseteq U$ of G' such that G'/L_U is a Lie group.*

We will also make use of the following lemma [9, Lemma 1.6.].

Lemma 8. *Let G be a locally compact group. Suppose that $N_1, N_2, \dots, N_k$ are closed normal subgroups, such that each for each i , G/N_i is a Lie group. Then $G/(N_1 \cap N_2 \cap \dots \cap N_k)$ is a Lie group.*

From now on unless otherwise stated φ_C always denotes the canonical projection from G onto G/C , where C is a compact normal subgroup.

Definition 9. *Suppose that G is a topological group. Then G_e denotes the connected component of the identity.*

Lemma 10. *Let G be a locally compact group, and $N, L \triangleleft G$ be compact normal subgroups, such that $G/N, G/L$ are Lie groups, and assume that G/N has λ -many connected components, i.e. $|G/N : (G/N)_e| = \lambda$. Then there is a finite number $k < \omega$, such that $G/(L \cap N)$ has $\leq k\lambda$ -many connected components.*

Proof. By Lemma 8, $G/(L \cap N)$ is a Lie group. Now, applying the second isomorphism theorem for topological groups (see [1, Thm. 1.5.18.]), $(G/(L \cap N)) / (L/(L \cap N)) \simeq G/L$, and $(G/(L \cap N)) / (N/(L \cap N)) \simeq G/N$ are Lie groups, thus replacing G by $G/(L \cap N)$, (and N, L by $N/(L \cap N), L/(L \cap N)$, respectively) we can assume that $L \cap N = \{e\}$.

Since N is a compact subgroup, by Cartan's theorem [12, Thm. 20.12] it is a Lie group too. By its compactness, N has only finitely many (say, n) connected components. We define $k = n$, and we will show that the open subgroup

$$H := \varphi_N^{-1}((G/N)_e) \leq G \text{ has at most } k \text{ connected components.} \quad (2)$$

($N \triangleleft H$, and $\varphi_N|_H : H \rightarrow (G/N)_e$ is a surjective open continuous homomorphism, thus $H/N \simeq (G/N)_e$.) Provided that (2) is true, it can be seen from the following that we are done. For any $\psi : G_1 \rightarrow G_2$ surjective homomorphism between the groups G_1, G_2 and $X \leq G_2$ subgroup

$$|G_2 : X| = |G_1 : \psi^{-1}(X)|,$$

since the preimage of the disjoint cosets $gX, g'X$ are the disjoint sets $\psi^{-1}(gX) = \psi^{-1}(g)\psi^{-1}(X)$, $\psi^{-1}(g'X) = \psi^{-1}(g')\psi^{-1}(X)$, which are cosets of $\psi^{-1}(X)$. Thus the index of H

$$|G : H| = |\varphi_N^{-1}(G/N) : \varphi_N^{-1}((G/N)_e)| = |G/N : (G/N)_e| = \lambda.$$

From which it clearly follows that the index $|G : H_e| = \lambda \cdot k$. But $H \leq G$ is an open subgroup of a Lie group, therefore H is the union of some cosets of G_e , and $G_e = H_e$, thus $|G : G_e| = \lambda \cdot k$.

Turning to the proof of (2), let φ'_N denote the restriction of φ_N to H , i.e. $\varphi'_N = \varphi_N|_H : H \rightarrow (G/N)_e$ is a surjective open continuous homomorphism. Recall that $H \leq G$ is open, thus H is a Lie group itself, thus H_e is open in H .

It is obvious that $N_e H_e$ is a connected open subset of H containing H_e (continuous image of the connected set $N_e \times H_e$). Therefore it is the component of the identity,

i.e. $N_e H_e = H_e$. This implies that if $N = \bigcup_{i < k} n_i N_e$, then

$$\bigcup_{i < k} n_i H_e = \bigcup_{i < k} n_i N_e H_e = \left(\bigcup_{i < k} n_i N_e \right) G_e = N H_e = (\varphi'_N)^{-1}(\varphi'_N(H_e)) \quad (3)$$

and that is a clopen subset of H (the union of left cosets of the open subgroup H_e), so is its complement. But $N H_e$ is the preimage of the set $\varphi_N(H_e) = \varphi_N(N H_e)$ under φ_N , thus its complement

$$H \setminus N H_e = (\varphi'^{-1}_N(H/N)) \setminus (\varphi'^{-1}_N(\varphi'_N(N H_e))) = \varphi'^{-1}_N((H/N) \setminus (\varphi_N(N H_e)))$$

is the preimage of $(H/N) \setminus (\varphi'_N(H_e))$. Therefore the clopen sets $N H_e$ and its complement is mapped by φ'_N onto the disjoint open sets $\varphi'_N(H_e) \subseteq H/N$ and its complement. But $H/N \simeq (G/N)_e$ is connected, thus at least one of these images is empty. Now $H_e \neq \emptyset$, thus $\varphi'_N(N H_e) \neq \emptyset$, which means that $G \setminus N H_e = \emptyset$, i.e. $G = N H_e$. From this $H = N H_e = \bigcup_{i < k} n_i H_e$, i.e. H is the union of (at most) k connected open subsets. $\square$

Corollary 11. *content...*

Now we can deduce the following.

Lemma 12. *Let G be a locally compact group. Then there is an open subgroup H , such that for every neighborhood $U \subseteq H$ of the identity, there is a compact normal subgroup $N_U \subseteq U$ of H , s.t. H/N_U is a Lie group that has finitely many connected components.*

Proof. Choose an arbitrary $G' \leq G$ which is given by Lemma 7, and let $K \triangleleft G'$ be a compact normal subgroup, such that G'/K is a Lie group. Let $\varphi_K : G' \rightarrow K$ denote the canonical projection. Then, since G'/K is a Lie group, $(G'/K)_e$ is open, thus $H := \varphi_K^{-1}((G'/K)_e)$ is open in G' . But G' is open in G , hence H is open in G .

First, we will check that H satisfies the conclusion of Lemma 7 (i.e. for every open set U ($e \in U \subseteq H$) there exists a compact normal subgroup $L_U \triangleleft H$, such that $L_U \subseteq U$, and H/L_U is a Lie group).

Let $U \subseteq H$ be an open neighborhood of the identity. Then there is a compact normal subgroup $L_U \triangleleft G'$, such that $L_U \subseteq U$ and G'/L_U is a Lie group. Then $L_U \triangleleft H \leq G'$ is obviously a normal subgroup of H . But if $\varphi_{L_U} : G' \rightarrow G'/L_U$ denotes the canonical projection, using that $H \subseteq G'$ is open, and φ_{L_U} is an open mapping, $\varphi_{L_U}(H) = H/L_U \leq G'/L_U$ is an open subgroup of a Lie group, thus H/L_U is a Lie group.

Let $N_U = K \cap L_U$. Then $H/(K \cap L_U)$ is a Lie group by Lemma 8, and since $H/K \simeq (G'/K)_e$ is connected, using Lemma 10 we obtain that $H/(K \cap L_U)$ has finitely many connected components. Thus choosing $N_U = K \cap L_U$ works. $\square$

Our plan is to take the subgroup H given by the previous lemma, and for an appropriately chosen compact normal subgroup $L \triangleleft H$ (where $L \leq K$), we will

construct a null but non-meager subgroup $S \leq H/L$. Then at first we would like to ensure that the preimage of S is also non-meager (in H), and hence it is also non-meager in G . That is what the following lemmas will be about.

We will need lemmas which are stated in terms of inverse limits.

For an arbitrary topological group G , let $(\mathfrak{C}, \preceq)$ denote the directed system of compact normal subgroups, with the reversed inclusion relation, that is,

$$\mathfrak{C} = \{C \triangleleft G : C \text{ is compact}\},$$

$$N \preceq M \iff N \geq M,$$

Now consider the following inverse system of topological groups.

$$(\{G/N : N \in \mathfrak{C}\}, \{\varphi_{M,N} : G/N \rightarrow (G/N)/(M/N) \simeq G/M : M \preceq N \in \mathfrak{C} \text{ (i.e. } N \leq M)\}).$$

Definition 13. Let G_a ($a \in I$) be an inverse system of topological groups $((I, \preceq)$ is a directed set), and let $J \subseteq I$ be a directed subset (wrt. $\preceq$). Then let $\pi_{I \rightarrow J}$ denote the natural projection

$$(x_a)_{a \in I} \mapsto (x_a)_{a \in J},$$

which is obviously a continuous homomorphism. Indeed, if we consider the projection on the whole product $\prod_{a \in I} G_a$, then it is continuous, and $\pi_{I \rightarrow J}$ is a restriction of it to the inverse limit. And clearly

$$\text{Ker}(\pi_{I \rightarrow J}) = (\varprojlim_{a \in I} G_a) \cap \left(\prod_{a \in J} \{e_{G_a}\} \times \prod_{a \in I \setminus J} G_a \right).$$

These projections may not be surjective.

Fix a locally compact group G , for which in every open neighborhood U of the identity there exists a compact normal subgroup N_U , such that G/N_U is a Lie group with finitely many connected components. Let $(\mathfrak{N}, \preceq)$ denote the following directed subsystem of the compact normal subgroups $\mathfrak{C}$

$$\mathfrak{N} = \{N \triangleleft G : N \in \mathfrak{C} \wedge G/N \text{ is a Lie group with finitely many connected components}\}$$

$$N \preceq M \iff N \geq M.$$

Notice that, by Lemmas 8, and 10 we obtain the following lemma which we will make use of later.

Lemma 14. Let G be a locally compact group such that in every open neighborhood U of the identity there exists a compact normal subgroup N_U , such that G/N_U is a Lie group with finitely many connected components. Then

$\mathfrak{N} = \{N \triangleleft G : N \triangleleft G \text{ compact} \wedge G/N \text{ is a Lie group with finitely many connected components}\}$ is directed wrt. the reversed inclusion relation.

Lemma 15. Let G be a topological group, and let $\mathfrak{M} \subseteq \mathfrak{C}$ be a directed subsystem of compact normal subgroups of G (i.e. for each $M, M' \in \mathfrak{M}$ there is an $M'' \in \mathfrak{M}$ with $M, M'' \geq M'$), and set $L := \bigcap_{M \in \mathfrak{M}} M$, then

- the mapping

$$\psi : G \rightarrow \varprojlim_{M \in \mathfrak{M}} G/M$$

$$g \mapsto \pi_{\mathfrak{M} \rightarrow \mathfrak{M}}((gN)_{N \in \mathfrak{N}}) = (gM)_{M \in \mathfrak{M}}$$

is a surjective open continuous homomorphism, and its kernel is L ,

- moreover the induced mapping $\bar{\psi} : G/L \rightarrow \varprojlim_{M \in \mathfrak{M}} G/M$ is an isomorphism.

Proof. First, ψ is obviously a continuous homomorphism.

- (i) For the surjectivity, let $(g_\alpha M_\alpha)_{M_\alpha \in \mathfrak{M}} \in \varprojlim_{M \in \mathfrak{M}} G/M$. Since the bonding maps $G/M \rightarrow G/M'$ ($M \leq M'$) are the canonical projections, the fact that $(g_\alpha M_\alpha)_{M_\alpha \in \mathfrak{M}}$ is in the inverse limit means that

$$\text{if } M_\alpha \geq M_\beta, \text{ then } g_\alpha M_\alpha \supseteq g_\beta M_\beta \quad (4)$$

From that it can be easily seen that the system $\{g_\alpha M_\alpha : M_\alpha \in \mathfrak{M}\} \subseteq \mathcal{P}(G)$ of compact sets has the finite intersection property.

By induction it is enough to show that for each pair $g_\alpha M_\alpha, g_\beta M_\beta$, there is a third element $g_\gamma M_\gamma$ in the system for which

$$g_\gamma M_\gamma \subseteq g_\alpha M_\alpha \cap g_\beta M_\beta.$$

But by the directedness, there is an $M_\gamma \in \mathfrak{M}$: $M_\gamma \leq M_\alpha, M_\beta$, then by (4):

$$g_\gamma M_\gamma \subseteq g_\alpha M_\alpha, g_\beta M_\beta.$$

Now, there is an element $g \in \bigcap_{M_\alpha \in \mathfrak{M}} g_\alpha M_\alpha \neq \emptyset$, then it can be easily seen that

$$\psi(g) = (gM_\alpha)_{M_\alpha \in \mathfrak{M}} = (g_\alpha M_\alpha)_{M_\alpha \in \mathfrak{M}}$$

(because $g \in g_\alpha M_\alpha$, i.e. $g = g_\alpha m_\alpha$ for some $m_\alpha \in M_\alpha$, thus $gM_\alpha = g_\alpha m_\alpha M_\alpha = g_\alpha M_\alpha$).

- (ii) The kernel of ψ is L indeed, since

$$Ker(\psi) = \psi^{-1}((M)_{M \in \mathfrak{M}}) = \bigcap_{M \in \mathfrak{M}} M = L.$$

- (iii) For the openness of ψ , since ψ is a homomorphism between topological groups, we only have to check that for each open set U containing the identity $\psi(U)$ is a neighborhood of the identity. Let U be an open set containing the identity, and, since $Im(U) = Im(UL)$ (recall that $Ker(\psi) = L$), we can assume that $UL = U$ (the set VX is open for arbitrary $X \subseteq G$ if $V \subseteq G$ is open). From $UL = U$ follows that $L \subseteq U$.

Now, using the compactness of the M -s, we will show that at least one of the M -s must be contained in U . Indeed, if the $M \setminus U$ -s are nonempty compact sets (for all $M \in \mathfrak{M}$), then because of the

directedness, the system $\{M \setminus U : M \in \mathfrak{M}\}$ of compact sets has the finite intersection property, thus

$$\emptyset \neq \bigcap_{M \in \mathfrak{M}} (M \setminus U) = \left(\bigcap_{M \in \mathfrak{M}} M \right) \setminus U = L \setminus U = \emptyset.$$

Thus there is an $M_0 \in \mathfrak{M}$ with $M_0 \subseteq U$. Then by some standard compactness argument, there is an open U' with $e \in U'$ such that $U'M_0 \subseteq U$.

Now we will check that the basic open set

$$B = \left(\varprojlim_{M \in \mathfrak{M}} G/M \right) \cap \left(\{xM_0 : x \in U'\} \times \prod_{M \in \mathfrak{M} \setminus \{M_0\}} G/M \right)$$

is contained in $\psi(U)$. Fix a $b \in B$. By the surjectivity of ψ , $b = \psi(g)$ for some $g \in G$

$$\pi_{\mathfrak{M} \rightarrow \{M_0\}}(\psi(g)) = gM_0 \in \{xM_0 : x \in U'\},$$

then $g \in gM_0 \subseteq U'M_0 \subseteq U$, i.e. $g \in U$, hence

$$b = \psi(g) \in \psi(U),$$

and

$$\psi(U) \supseteq B$$

thus ψ is an open mapping.

- For our second claim, observe that for any open subset $U \subseteq G/L$

$$\bar{\psi}(U) = \psi(\varphi_L^{-1}(U))$$

is open by openness of ψ , and the continuity of the canonical projection $\varphi_L : G \rightarrow G/L$.

□

Corollary 16. *Let G be a locally compact group, such that for every open neighborhood U of the identity there exists a compact normal subgroup $N_U \subseteq U$, for which G/N_U is a Lie group having finitely many connected components.*

(i) *Then the mapping*

$$g \mapsto (gN)_{N \in \mathfrak{N}}$$

is an isomorphism, hence

$$G \simeq \varprojlim_{N \in \mathfrak{N}} G/N.$$

(ii) *For any directed subsystem $\mathfrak{M}$,*

$$\text{the projection } \pi_{\mathfrak{N} \rightarrow \mathfrak{M}} \text{ is a surjective open mapping.} \quad (5)$$

Proof. (i) We only have to check the conditions of Lemma 15:

- $\mathfrak{N} = \{N \triangleleft G : N \text{ is compact, } G/N \text{ is a Lie group with finitely many connected components}\}$ is directed, but (14) stated that this is true.

- $\bigcap_{N \in \mathfrak{N}} N = \{e\}$:
For any $g \in G$, if $g \neq e$, then there is an open neighborhood U of the identity such that $g \notin U$. Then G/N_U is a Lie group with finitely many components, i.e. $N_U \in \mathfrak{N}$.
- (ii) Our second claim easily follows from that $\pi_{\mathfrak{N} \rightarrow \mathfrak{M}}$ is the composition of the inverse of the homeomorphism: $\varphi : G \rightarrow \varprojlim_{N \in \mathfrak{N}} G/N$ ($g \mapsto (gN)_{N \in \mathfrak{N}}$), and the aforementioned open projection $\psi_{\mathfrak{M}} : G \rightarrow \varprojlim_{M \in \mathfrak{M}} G/M$ ($g \mapsto (gM)_{M \in \mathfrak{M}}$):

$$\pi_{\mathfrak{N} \rightarrow \mathfrak{M}} = \psi_{\mathfrak{M}} \circ \varphi^{-1}.$$

□

Now, we can turn to a key lemma:

Lemma 17. *Let $(\{G_\alpha : \alpha \in I\}, \{\varphi_{\alpha, \beta} : \alpha \preceq \beta \in I\})$ be an inverse system of second countable topological groups.*

Assume that a countable subset $J_0 \subseteq I$ of the indexing poset is given, and let V_k ($k \in \omega$) be dense open sets in $\varprojlim_{\alpha \in I} G_\alpha$. Assume that for any directed subsystem $H \subseteq I$ the projection

$$\pi_{I \rightarrow H} : \varprojlim_{i \in I} G_i \rightarrow \varprojlim_{i \in H} G_i \text{ is surjective.} \quad (6)$$

Then there is a countable directed subsystem $J \subseteq I$, $J_0 \subseteq J$, and a sequence $(V'_k)_{k \in \omega}$ of dense open sets in $\varprojlim_{\alpha \in J} G_\alpha$, such that for each $k \in \omega$

$$\pi_{I \rightarrow J}^{-1}(V'_k) \subseteq V_k.$$

Remark 18. *Condition (6) could be easily dropped, but in our applications this condition will always hold, thus for simplicity, and for avoiding confusions we assume it.*

Proof. We can assume that $J_0 \neq \emptyset$ (by choosing an arbitrary $\alpha \in I$ to be an element of J_0). We will find a countable set J such that: $J_0 \subseteq J \subseteq I$, and which is directed, and closed under the following countably many operations. For each $\alpha \in I$ fix a countable base $\mathcal{B}_\alpha = \{B_1^\alpha, B_2^\alpha, \dots\}$ consisting of nonempty open subsets of G_α . Recall that there is a base of $\varprojlim_{\alpha \in I} G_\alpha$ consisting of sets of the form

$$D_n^\alpha := \left(B_n^\alpha \times \prod_{\beta \in I \setminus \{\alpha\}} G_\beta \right) \cap \varprojlim_{\gamma \in I} G_\gamma \quad (n \in \omega, \alpha \in I).$$

(Since B_n^α -s are nonempty open sets, by (6) the D_n^α -s are also nonempty.) Now for every $n, m \in \omega$ we define the following operation, which is a choice function.

$$f_{n,m} : I \rightarrow I$$

$$\alpha \mapsto \beta \quad \text{s.t.} \quad \exists k : D_k^\beta \subseteq V_n \cap D_m^\alpha$$

(such a β exists because each V_n is a dense open set). And let $g : I \times I \rightarrow I$ be another choice function such that $\alpha, \beta \preceq g(\alpha, \beta)$, i.e. a subset of I closed under the

function g is a directed subsystem. Then, take the closure J of J_0 under the $f_{n,m}$ -s and g . Then, because J is closed under g , it is a directed system.

Similarly to the D_n^β -s, we define nonempty basic open sets in the limit $\varprojlim_{\gamma \in J} G_\gamma$ as follows.

$$E_n^\alpha := \left(B_n^\alpha \times \prod_{\beta \in J \setminus \{\alpha\}} G_\beta \right) \cap \varprojlim_{\gamma \in J} G_\gamma \quad (n \in \omega, \alpha \in J).$$

We have the directed system $J \subseteq I$, and for each $n, m \in \omega$, and $\alpha \in J$, since J is closed under the function $f_{n,m}$, there is a $k = k_{n,m,\alpha} \in \omega$, and $\beta = \beta_{n,m,\alpha} \in J$ such that

$$D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} \subseteq V_n \cap D_m^\alpha \quad (7)$$

Next we claim that for each $n, m \in \omega$, and $\alpha \in J$, if $k_{n,m,\alpha} \in \omega$, and $\beta_{n,m,\alpha} \in J$ are what (7) gives, then

$$E_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} \subseteq E_m^\alpha \quad (8)$$

and

$$D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} = \pi_{I \rightarrow J}^{-1}(E_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}}) \subseteq V_n. \quad (9)$$

For (8), first let $k = k_{n,m,\alpha} \in \omega$, and $\beta = \beta_{n,m,\alpha} \in J$. Reformulating the condition $D_k^\beta \subseteq D_m^\alpha$ (following from (7)), we get that

$$\left[(x_\gamma)_{\gamma \in I} \in \varprojlim_{\gamma \in I} G_\gamma \wedge x_\beta \in B_k^\beta \right] \Rightarrow (x_\gamma)_{\gamma \in I} \in D_m^\alpha \quad (\iff x_\alpha \in B_m^\alpha).$$

Now notice that by our assumptions, the projection $\pi_{I \rightarrow J} : \varprojlim_{\gamma \in I} G_\gamma \rightarrow \varprojlim_{\gamma \in J} G_\gamma$ is surjective, thus for a fixed element $(x_\gamma)_{\gamma \in J}$ in the inverse limit of the smaller system there is an element $(y_\gamma)_{\gamma \in I} \in \varprojlim_{\gamma \in I} G_\gamma$ with $(x_\gamma)_{\gamma \in J} = \pi_{I \rightarrow J}((y_\gamma)_{\gamma \in I})$. Then

$$\begin{aligned} (x_\gamma)_{\gamma \in J} \in E_k^\beta &\iff \left[(x_\gamma)_{\gamma \in J} \in \varprojlim_{\gamma \in J} G_\gamma \wedge x_\beta \in B_k^\beta \right] \iff \\ &\iff \left[(x_\gamma)_{\gamma \in J} = \pi_{I \rightarrow J}((y_\gamma)_{\gamma \in I}), (y_\gamma)_{\gamma \in I} \in \varprojlim_{\gamma \in I} G_\gamma \wedge y_\beta = x_\beta \in B_k^\beta \right] \Rightarrow \\ &\Rightarrow [y_\alpha = x_\alpha \in B_m^\alpha] \iff (x_\gamma)_{\gamma \in J} \in E_m^\alpha, (y_\gamma)_{\gamma \in I} \in D_m^\alpha. \end{aligned}$$

This clearly yields (8), as desired.

For (9), $D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} \subseteq V_n$ is true by (7), we only have to check that $D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} = \pi_{I \rightarrow J}^{-1}(E_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}})$. But for arbitrary $l \in \omega$ and $\delta \in J$

$$D_l^\delta = \pi_{I \rightarrow J}^{-1}(E_l^\delta),$$

since for $(x_\gamma)_{\gamma \in I} \in \varprojlim_{i \in I} G_i$

$$\begin{aligned} (x_\gamma)_{\gamma \in I} \in D_l^\delta &\iff x_\delta \in B_l^\delta \iff \\ &\iff (x_\gamma)_{\gamma \in J} = \pi_{I \rightarrow J}((x_\gamma)_{\gamma \in I}) \in E_l^\delta \end{aligned}$$

Finally, set $V'_n = \bigcup_{m \in \omega, \alpha \in J} E_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}}$. Since m ranges over ω and α over J , using (8) V'_n is a dense open set. From (9)

$$\pi_{I \rightarrow J}^{-1}(V'_n) = \bigcup_{m \in \omega, \alpha \in J} \pi_{I \rightarrow J}^{-1}(E_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}}) = \bigcup_{m \in \omega, \alpha \in J} D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}},$$

and using (7),

$$\bigcup_{m \in \omega, \alpha \in J} D_{\beta_{n,m,\alpha}}^{k_{n,m,\alpha}} \subseteq V_n.$$

□

This lemma has the following consequence.

Lemma 19. *Assume that G is a locally compact group, in which for every open neighborhood U of the identity there is a compact normal subgroup N_U , such that G/N_U is a Lie group with finitely many connected components. Let a countable set $J_0 = \{N_{U_i} : i \in \omega\} \subseteq \mathfrak{N}$ be given. Assume that $R \subseteq G$ is co-meager. Then there exist a countable set $J \supseteq J_0$, $J \subseteq \mathfrak{N}$, and a compact normal subgroup $K' = \bigcap_{N \in J} N$ of G , and a co-meager set $R' \subseteq G/K'$, such that $\varphi_{K'}^{-1}(R') \subseteq R$, and G/K' is an inverse limit of countable many second countable Lie groups, thus is a Polish group.*

Proof. Since, by Corollary 16, $G \simeq \varprojlim_{N \in \mathfrak{N}} G/N$, where $\varphi : G \rightarrow \varprojlim_{N \in \mathfrak{N}} G/N$ denotes the canonical isomorphism (i.e. $g \mapsto (gN)_{N \in \mathfrak{N}}$).

Recall that, by Lemma 15, for each directed subsystem $\mathfrak{M} \subseteq \mathfrak{N}$, the projection $\pi_{\mathfrak{N} \rightarrow \mathfrak{M}}$ is also surjective, and open.

First, we will work in $\varprojlim_{N \in \mathfrak{N}} G/N$.

Let $\hat{R} = \varphi(R)$ denote the corresponding co-meager subset of $\varprojlim_{N \in \mathfrak{N}} G/N$. We know that the G/N -s ($N \in \mathfrak{N}$) are Lie groups having finitely many connected components, thus are second countable.

Let $(V_m)_{m \in \omega}$ be a sequence of dense open sets in $\varprojlim_{N \in \mathfrak{N}} G/N$ such that $\bigcap_{m \in \omega} V_m \subseteq \hat{R}$. We will apply Lemma 17 with $I = \mathfrak{N}$, the sequence $(V_m)_{m \in \omega}$, we only have to check (6). Recall that, by Lemma 15, for each directed subsystem $\mathfrak{M} \subseteq \mathfrak{N}$, the projection $\pi_{\mathfrak{N} \rightarrow \mathfrak{M}}$ is surjective.

Applying Lemma 17, there is a countable directed subset $J \subseteq \mathfrak{N}$ ($J_0 \subseteq J$), and a sequence $(V'_m)_{m \in \omega}$ of dense open sets in $\varprojlim_{N \in J} G/N$ such that

$$\pi_{\mathfrak{N} \rightarrow J}^{-1}(V'_m) \subseteq V_m \quad (\forall m \in \omega).$$

Hence, if we set the co-meager set $\hat{R}' := \bigcap_{m \in \omega} V'_m$ then clearly

$$\pi_{\mathfrak{N} \rightarrow J}^{-1}(\hat{R}') = \pi_{\mathfrak{N} \rightarrow J}^{-1} \left(\bigcap_{m \in \omega} V'_m \right) = \bigcap_{m \in \omega} \pi_{\mathfrak{N} \rightarrow J}^{-1}(V'_m) \subseteq \bigcap_{m \in \omega} V_m \subseteq \hat{R}.$$

But $\varprojlim_{N \in J} G/N$ is a quotient of $\varprojlim_{N \in \mathfrak{N}} G/N$, since we saw that the projection $\pi_{\mathfrak{N} \rightarrow J}$ is surjective and open. Thus, by the first isomorphism theorem for topological groups (see [1, Lemma 1.5.13.]) $\varprojlim_{N \in J} G/N$ is isomorphic to the topological group $(\varprojlim_{N \in \mathfrak{N}} G/N) / \text{Ker}(\pi_{\mathfrak{N} \rightarrow J})$, i.e. it is a quotient of $(\varprojlim_{N \in \mathfrak{N}} G/N)$.

Now we have a co-meager set $\hat{R}'$ with $\pi_{\mathfrak{N} \rightarrow J}^{-1}(\hat{R}') \subseteq \hat{R}$ in

$$\varprojlim_{N \in J} G/N \simeq (\varprojlim_{N \in \mathfrak{N}} G/N) / \text{Ker}(\pi_{\mathfrak{N} \rightarrow J}).$$

Let $L := \text{Ker}(\pi_{\mathfrak{N} \rightarrow J})$, and

$$\varphi_L : \left(\varprojlim_{N \in \mathfrak{N}} G/N \right) \rightarrow \left(\varprojlim_{N \in \mathfrak{N}} G/N \right) / L$$

denote the projection, let $\bar{\psi} : \left(\varprojlim_{N \in \mathfrak{N}} G/N \right) / L \rightarrow \varprojlim_{N \in J} G/N$ denote the isomorphism for which

$$\pi_{\mathfrak{N} \rightarrow J} = \bar{\psi} \circ \varphi_L. \quad (10)$$

Set $\hat{R}'' = \bar{\psi}^{-1}(\hat{R}')$, by (10) that is a co-meager set such that $\varphi_L^{-1}(\hat{R}'') \subseteq \hat{R}$. Now we have a normal subgroup $L = \text{Ker}(\pi_{\mathfrak{N} \rightarrow J})$ in $\varprojlim_{N \in \mathfrak{N}} G/N$, such that there is a co-meager $\hat{R}''$ in the quotient $\left(\varprojlim_{N \in \mathfrak{N}} G/N \right) / L$, where

$$\varphi_L^{-1}(\hat{R}'') \subseteq \hat{R} = \varphi(R). \quad (11)$$

We only have to pull back this construction to G by φ , and check that the normal subgroup $L = \text{Ker}(\pi_{\mathfrak{N} \rightarrow J})$ is a compact subgroup, or equivalently its pull-back $\varphi^{-1}(\text{Ker}(\pi_{\mathfrak{N} \rightarrow J}))$ is compact.

Since

$$\text{Ker}(\pi_{\mathfrak{N} \rightarrow J}) = \left(\prod_{N \in J} \{N\} \times \prod_{N \in \mathfrak{N} \setminus J} G/N \right) \cap \varprojlim_{N \in \mathfrak{N}} G/N,$$

we have that

$$\varphi^{-1}(\text{Ker}(\pi_{\mathfrak{N} \rightarrow J})) = \varphi^{-1} \left(\prod_{N \in J} \{N\} \times \prod_{N \in \mathfrak{N} \setminus J} G/N \right) = \bigcap_{N \in J} N,$$

that is indeed a compact subgroup.

Let $K' = \varphi^{-1}(\text{Ker}(\pi_{\mathfrak{N} \rightarrow J}))$. Now, since $\varphi : G \rightarrow \varprojlim_{N \in \mathfrak{N}} G/N$ is an isomorphism, which maps K' onto $L = \text{Ker}(\pi_{\mathfrak{N} \rightarrow J})$, there is an induced isomorphism $\bar{\varphi} : G/K' \rightarrow \left(\varprojlim_{N \in \mathfrak{N}} G/N \right) / L$. Now it is easy to check that if we set $R' = \bar{\varphi}^{-1}(\hat{R}'')$, then by (11)

$$\varphi_{K'}(R') \subseteq R.$$

Finally, it remains to check that

$$G/K' \simeq \left(\varprojlim_{N \in \mathfrak{N}} G/N \right) / \text{Ker}(\pi_{\mathfrak{N} \rightarrow J}) \simeq \varprojlim_{N \in J} G/N$$

is an inverse limit of countably many Polish spaces. We know that $\varprojlim_{N \in J} G/N$ is an inverse limit of countably many Lie groups, each having finitely many connected components, from which each G/N is a Polish group. Therefore $\varprojlim_{N \in J} G/N$ is an inverse limit of countably many Polish groups, i.e. it is a closed subset of the product of countably many Polish groups, thus is Polish.

□

Lemma 20. *Assume that G is a locally compact group, in which for every open neighborhood U of the identity there is a compact normal subgroup N_U , such that G/N_U is a Lie group with finitely many connected components. Let a countable set*

$J_0 = \{N_{U_i} : i \in \omega\} \subseteq \mathfrak{N}$ be given. Set $K := \bigcap_{N \in J_0} N$, and let $M \subseteq G/K$ be a non-meager subset (in the factor topology). Then $\varphi_K^{-1}(M)$ is non-meager in G .

Proof. On the contrary, assume that the set $R = G \setminus \varphi_K^{-1}(M)$ is co-meager. Then (by Lemma 19) there is a compact normal subgroup $K' \leq K$, and co-meager set $R' \subseteq G/K'$, such that G/K' is Polish, and

$$\varphi_{K'}^{-1}(R') \subseteq R,$$

i.e.

$$\varphi_{K'}^{-1}(R') \cap \varphi_K^{-1}(M) = \emptyset.$$

Then clearly (in G/K')

$$R' \cap \varphi_{K/K'}^{-1}(M) = \emptyset. \quad (12)$$

But

$$\varphi_{K/K'} : G/K' \rightarrow G/K$$

is a surjective open mapping between Polish spaces (since by Lemma 15, $G/K \simeq \varprojlim_{N \in J_0} G/N$ is the inverse limit of countably many Polish spaces, thus is Polish), and according to [2, Lemma 2.6] it maps a co-meager set onto a co-meager set. Thus $\varphi_{K/K'}(R') \subseteq G/K$ is co-meager. If we could check that

$$\varphi_{K/K'}(R') \cap M = \emptyset$$

then this would be a contradiction, since M was of second category. But using (12) and $G/K' = \varphi_{K/K'}^{-1}(G/K)$

$$R' \subseteq (G/K') \setminus \varphi_{K/K'}^{-1}(M) = \varphi_{K/K'}^{-1}((G/K) \setminus M),$$

thus $\varphi_{K/K'}(R') \subseteq (G/K) \setminus M$, as desired. $\square$

We will need that if $H \leq G$ is an open subgroup, $S \leq H/L$ (with some compact $L \triangleleft H$) is null with respect to the Haar measure of H/L , then $\varphi_K^{-1}(S)$ is null in H . (That would imply that it is null in G , since for an open subgroup H the restriction of the Haar measure $\mu_G|_H$ is a Haar measure of H .)

The following lemma can be found [10, Sec.63, Theorem C].

Lemma 21. *Let G be a locally compact group, and μ be a left-invariant Borel measure on it, which is positive on nonempty open sets, and finite on compact sets, and let $K \triangleleft G$ be a compact normal subgroup. Then the push-forward measure $B \mapsto \mu(\varphi_K^{-1}(B))$ is a left-invariant Borel measure on G/K , which is positive on nonempty open sets, and finite on compact sets.*

The following lemma is well-known, but for the sake of completeness we include a proof:

Lemma 22. *Let G be a locally compact group, and*

$$\mu : \mathcal{B}(G) \rightarrow [0, \infty]$$

a left Haar Borel measure on it and let $K \triangleleft G$ be a compact normal subgroup, ν a left Haar Borel measure on G/K . Then ν is the same (up to a multiplicative

constant) as the push-forward of μ , i.e. there is a constant $c > 0$ such that for all $H \subseteq G/K$ Borel

$$c\nu(H) = \mu(\varphi_K^{-1}(H)).$$

Proof. By Lemma 21, the push-forward of μ is a left-invariant Borel measure on G/K , which is positive on nonempty open sets, and finite on compact sets. By the uniqueness, we only have to check the regularity. Let $B \subseteq G/K$ be a Borel subset, then, since taking the supremum is monotonic, and continuous image of a compact set is compact,

$$\begin{aligned} \sup\{\mu(\varphi_K^{-1}(C)) : C \subseteq B \subseteq G/K, C \text{ is compact}\} &\geq \\ \sup\{\mu(\varphi_K^{-1}(\varphi(D))) : D \subseteq \varphi_K^{-1}(B) \subseteq G \text{ is compact}\} &= \\ \sup\{\mu(DK) : D \subseteq \varphi_K^{-1}(B) \subseteq G \text{ is compact}\} &\geq \\ \sup\{\mu(D) : D \subseteq \varphi_K^{-1}(B) \subseteq G \text{ is compact}\}. \end{aligned}$$

(The first inequality holds, because if $D \subseteq \varphi_K^{-1}(B)$ is compact, then $\varphi_K(D) \subseteq B$ is compact, and the equality is by $\varphi_K^{-1}(\varphi_K(D)) = DK$.) And

$$\sup\{\mu(D) : D \subseteq \varphi_K^{-1}(B) \subseteq G \text{ is compact}\} = \mu(\varphi_K^{-1}(B))$$

by the regularity of μ . $\square$

Recall that under a left Haar measure we mean the completion of a left Haar Borel measure.

Corollary 23. *Let G be a locally compact group, μ is a left Haar measure on G . Suppose that $K \triangleleft G$ is a compact normal subgroup, ν is a left Haar measure on G/K . If $X \subseteq G/K$ is null wrt. ν , then $\varphi_K^{-1}(X)$ is null wrt. μ .*

Proof. Let $B \supseteq X$ be a null Borel set in G/K . Then, applying Lemma 22 for $\mu|_{\mathcal{B}(G)}$, and $\nu|_{\mathcal{B}(G/K)}$ we obtain that (by the continuity of φ_K) $\mu|_{\mathcal{B}(G)}(\varphi_K^{-1}(B)) = 0$, hence

$$\mu(\varphi_K^{-1}(X)) \leq \mu(\varphi_K^{-1}(B)) = 0.$$

$\square$

Before the construction for profinite groups, and connected Lie groups, we will need the following technical lemma about co-meager sets in compact metric spaces (the lemma uses the idea of the well known characterization of co-meager sets in the Cantor space, see [3, Thm. 2.2.4]).

Before stating the lemma, recall that in a metric space (X, d) , the diameter of a subset $S \subseteq X$ is defined as follows

$$\text{diam}(S) = \sup\{d(s, s') : s, s' \in S\}.$$

Lemma 24. *Assume that (X, d) is a compact metric space, and let $(m_i)_{i \in \omega}$ be a sequence of positive integers, let $R \subseteq X$ be a co-meager set.*

Assume that the system of compact sets $(C_{p_0 p_1 \dots p_i})_{p_0 p_1 p_2 \dots p_i \in \prod_{j=0}^i \{1, 2, \dots, m_j\}, i \in \omega}$ fulfills the following conditions:

- (1) $|\bigcap_{i \in \omega} C_{p_0 p_1 \dots p_i}| = 1$ for each $p_0 p_1 \dots p_m \dots \in \prod_{j \in \omega} \{1, 2, \dots, m_j\}$,
- (2) $C_{p_0 p_1 \dots p_n} \subseteq C_{p_0 p_1 \dots p_n p_{n+1}}$ for each $n \geq 0$ and $C_{p_0 p_1 \dots p_n p_{n+1}}$,
- (3) $\text{int}(C_{p_0 p_1 \dots p_i}) \neq \emptyset$ for each $C_{p_0 p_1 \dots p_i}$,
- (4) for each $C_{p_0 p_1 \dots p_i}$

$$C_{p_0 p_1 \dots p_i} = \bigcup_{1 \leq j \leq m_{i+1}} \overline{\text{int}(C_{p_0 p_1 \dots p_i j})}.$$

Then there is an increasing sequence $0 = n_0 < n_1 < \dots < n_j < \dots$ of integers, and an infinite sequence

$$r := (r_j)_{j \in \omega} \in \prod_{j \in \omega} \{1, 2, \dots, m_j\}$$

such that for each sequence $s = (s_j)_{j \in \omega} \in \prod_{j \in \omega} \{1, 2, \dots, m_j\}$, if $\{j \in \omega : s|_{[n_j, n_{j+1})} = r|_{[n_j, n_{j+1})}\}$ is infinite, then $\bigcap_{i \in \omega} C_{s_0 s_1 \dots s_i} \subseteq R$.

Proof. Let $\bigcap_{k \in \omega} U_k$ be the intersection of dense open sets such that $\bigcap_{k \in \omega} U_k \subseteq R$. By replacing U_k with $U_0 \cap U_1 \cap \dots \cap U_k$ we can assume that $(U_k)_{k \in \omega}$ is a decreasing sequence. We construct the sequences $(n_i)_{i \in \omega}$ and $(r_i)_{i \in \omega}$ simultaneously by induction on i as follows. Before the $i+1$ -th step the initial segments $0 = n_0 < n_1 < n_2 < \dots < n_i$, and $r_0 r_1 \dots r_{n_i-1}$ are already defined, and we will choose n_{i+1} , and $r_{n_i} r_{n_i+1} \dots r_{n_{i+1}-1}$ such that

$$\begin{aligned} C_{p_0 p_1 \dots p_{n_i-1} r_{n_i} r_{n_i+1} r_{n_i+2} \dots r_{n_{i+1}-1}} &\subseteq U_{i+1} \\ \text{for each } p_0 p_1 \dots p_{n_i-1} &\in \prod_{j < n_i} \{1, 2, \dots, m_j\}. \end{aligned} \quad (13)$$

First, we show that we can choose n_{i+1} and $r_{n_i} r_{n_i+1} \dots r_{n_{i+1}-1}$ so that (13) holds.

Assume that $n_0, n_1, n_2, \dots, n_i$ are defined. Now enumerate the set

$$\{C_{p_0 p_1 \dots p_{n_i-1}} : p_0 p_1 \dots p_{n_i-1} \in \prod_{j < n_i} \{1, 2, \dots, m_j\}\} = \{D_1, D_2, \dots, D_{m_0 m_1 m_2 \dots m_{n_i-1}}\}.$$

For each D_l define the sequences $\underline{r}^l = r_1^l r_2^l \dots r_{n_i}^l$ by induction on l , such that if $D_l = C_{p_0 p_1 \dots p_{n_i-1}}$, then for the concatenation $\underline{r}^1 \underline{r}^2 \dots \underline{r}^l$ of the $\underline{r}^j$ -s ($j \leq l$)

$$C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^l} \subseteq U_{i+1}. \quad (14)$$

Before constructing the $\underline{r}^j$ -s ($j \leq m_0 m_1 m_2 \dots m_{n_i-1}$), we prove that if we could manage to choose such $\underline{r}^l$ -s, then setting

- $n_{i+1} = n_i + \text{length}(\underline{r}^1 \underline{r}^2 \dots \underline{r}^{m_0 m_1 m_2 \dots m_{n_i-1}})$, and
- $r_{n_i} r_{n_i+1} \dots r_{n_{i+1}-1} = \underline{r}^1 \underline{r}^2 \dots \underline{r}^{m_0 m_1 m_2 \dots m_{n_i-1}}$

would yield (13). Since for $D_l = C_{p_0 p_1 \dots p_{n_i-1}}$ (2), and (14) implies that

$$C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{m_0 m_1 m_2 \dots m_{n_i-1}}} \subseteq C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^l} \subseteq U_{i+1},$$

i.e. (13) holds indeed.

Turning to the construction of the $\underline{r}^l$ -s, assume that $\underline{r}^1, \underline{r}^2, \dots, \underline{r}^{l-1}$ are defined. If $D_l = C_{p_0 p_1 \dots p_{n_i-1}}$, then, because

$$\text{int}(C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1}}) \neq \emptyset$$

and U_{i+1} is a dense open set, there is an $x \in X$, and an open ball

$$B(x, r) = \{y \in X : d(x, y) < r\} \subseteq C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1}} \cap U_{i+1}.$$

Now we can use (4), thus there is an r_0^l such that for the open set $V_0 = \text{int}(C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l})$

$$V_0 \cap B\left(x, \frac{r}{2}\right) \neq \emptyset.$$

Similarly, by (4), there are infinite sequences $r_0^l, r_1^l, r_2^l, \dots, r_k^l, \dots$ and $V_0, V_1, \dots$, where $V_k = \text{int}(C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l r_1^l \dots r_k^l})$ such that

$$V_k \cap B\left(x, \frac{r}{2}\right) \neq \emptyset.$$

And using (1) we know that

$$\lim_{k \rightarrow \infty} \text{diam}(C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l r_1^l \dots r_k^l}) = 0,$$

thus there is an index k_0 such that

$$\text{diam}(V_{k_0}) \leq \text{diam}(C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l r_1^l \dots r_{k_0}^l}) \leq \frac{r}{4},$$

from which

$$C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l r_1^l \dots r_{k_0}^l} \subseteq \overline{B\left(x, \frac{3r}{4}\right)},$$

thus

$$C_{p_0 p_1 \dots p_{n_i-1} \underline{r}^1 \underline{r}^2 \dots \underline{r}^{l-1} r_0^l r_1^l \dots r_{k_0}^l} \subseteq B(x, r) \subseteq U_{i+1}.$$

This finishes the construction of $(n_i)_{i \in \omega}$ and $(r_i)_{i \in \omega}$.

Now, as the r_i -s, and n_i -s are already constructed, we only have to check that for any sequence $(s_j)_{j \in \omega} \in \prod_{j \in \omega} \{1, 2, \dots, m_j\}$ if $\{j \in \omega : r_{[n_j, n_{j+1}]} = s_{[n_j, n_{j+1}]}\}$ is infinite, then $\bigcap_{i \in \omega} C_{s_0 s_1 \dots s_i} \subseteq R$. Fix such a sequence $s = (s_j)_{j \in \omega}$.

If $k \in \{j \in \omega : r_{[n_j, n_{j+1}]} = s_{[n_j, n_{j+1}]}\}$, then, using (13):

$$C_{s_0 s_1 \dots s_{n_k-1} r_{n_k} r_{n_k+1} r_{n_k+2} \dots r_{n_{k+1}-1}} \subseteq U_{k+1}.$$

Thus, if $x \in \bigcap_{i \in \omega} C_{s_0 s_1 \dots s_i}$, $k \in \{j \in \omega : r_{[n_j, n_{j+1}]} = s_{[n_j, n_{j+1}]}\}$ are fixed, then

$$x \in C_{s_0 s_1 \dots s_{n_{k+1}-1}} = C_{s_0 s_1 \dots s_{n_k-1} r_{n_k} r_{n_k+1} r_{n_k+2} \dots r_{n_{k+1}-1}} \subseteq U_{k+1},$$

hence $x \in U_{k+1}$ for infinitely many k -s. But recall that the U_k -s form a decreasing sequence, thus

$$x \in \bigcap_{k \in \omega} U_k \subseteq R.$$

□

Proposition 25. *Let G be the inverse limit of the following inverse system, consisting of finite groups.*

$$\{G_i, \varphi_{i,j} : G_j \rightarrow G_i, i \leq j \in \omega\},$$

i.e.

$$G = \{(g_i)_{i \in \omega} : \varphi_{j,j+1}(g_{j+1}) = g_j \ (\forall j)\} \leq \prod_{j \in \omega} G_j,$$

where each $|G_{j+1}|/|G_j| > 1$. Then there exists a subgroup $S \leq G$ that is null but non-meager.

Proof. Let $\varphi_i : G \rightarrow G_i$ denote the canonical projection. Since, by [13, Lemma 1.1.9], for an arbitrary increasing sequence $n_1 < n_2 < \dots$ of positive integers

$$\varprojlim_{i \in \omega} G_i \simeq \varprojlim_{i \in \omega} G_{n_i},$$

we can assume that the sequence

$$(m_i = \frac{|G_i|}{|G_{i-1}|})_{1 \leq i \in \omega}$$

can grow as rapidly as we want. Later we will recursively thin out the sequence of G_i -s. Another corollary of this lemma is that we can assume that $G_0 = \{e\}$ is the trivial group. Define $m_0 = 1$, thus from now $\text{dom}(m) = \omega$.

For each $g \in G_{i-1}$ ($i \geq 1$) fix an enumeration of the set

$$\varphi_{i-1,i}^{-1}(g) = \{g^{(1)}, g^{(2)}, \dots, g^{(m_i)}\},$$

so that

$$e_{G_{i-1}}^{(1)} = e_{G_i} \quad (15)$$

holds. Let $G_i^{(j)} \subseteq G_i$ denote the set $\{g^{(j)} : g \in G_{i-1}\}$, i.e. for each $g \in G_{i-1}$ it contains exactly one element from $\varphi_{i-1,i}^{-1}(g)$, we have a partition of G_i .

Fixing these enumerations, each element of G can be uniquely identified with an element of $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$ in the following way. Let $\psi_i : G_i \rightarrow \{1, 2, \dots, m_i\}$ denote the mapping

$$g \mapsto k \text{ iff } g \in G_i^{(k)}, \quad (16)$$

i.e. $g = (\varphi_{i-1,i}(g))^{(k)}$. Then consider the product $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$ with the product topology, and the mapping

$$\psi : \varprojlim_{i \in \omega} G_i \rightarrow \prod_{i \in \omega} \{1, 2, \dots, m_i\}$$

$$(g_i)_{i \in \omega} \mapsto (\psi_i(g_i))_{i \in \omega}.$$

Notice that for $e_G = (e_{G_i})_{i \in \omega} \in G$ by (15)

$$\psi(e_G) = (1)_{i \in \omega} \quad (17)$$

Claim 26. *The mapping $\psi : G \rightarrow \prod_{j \in \omega} \{1, 2, \dots, m_j\}$ is a homeomorphism. Furthermore, considering G with the probability left Haar measure μ_G (i.e. $\mu_G(G) = 1$), and the product space with the canonical measure ν , i.e.*

$$\nu(\{k_0\} \times \{k_1\} \times \dots \times \{k_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\}) = \frac{1}{m_0} \cdot \frac{1}{m_1} \cdot \dots \cdot \frac{1}{m_i},$$

ψ is measure preserving.

Proof. It is straightforward to check that ψ is a bijection.

Fix an element $g \in G_i$ of the i -th group and the corresponding basic set

$$B = \left(\{g\} \times \prod_{j \in \omega, j \neq i} G_j \right) \cap \varprojlim_{j \in \omega} G_j$$

in $\varprojlim_{j \in \omega} G_j$.

Then consider the following sequence

$$h_i = g, h_{i-1} = \varphi_{i-1,i}(g), \dots, h_0 = \varphi_{0,i}(g)$$

from which we can define k_j -s ($j \leq i, 1 \leq k_j \leq m_j$) such that $h_j = h_{j-1}^{(k_j)}$. Then one can easily see that

$$\psi(B) = \{k_0\} \times \{k_1\} \times \dots \times \{k_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\}.$$

Conversely, there is a base $\mathcal{B}$ of $\prod_{j \in \omega} \{1, 2, \dots, m_j\}$ containing sets of the form

$$\{l_0\} \times \{l_1\} \times \{l_2\} \times \dots \times \{l_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\}.$$

Fixing a basic open set

$$\{l_0\} \times \{l_1\} \times \{l_2\} \times \dots \times \{l_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\} \in \mathcal{B},$$

we extend recursively the sequence $g_0 = e_{G_0}$ of length 1 to $(g_j)_{j \leq i}$ such that $g_j = g_{j-1}^{(l_j)}$ for all $1 \leq j \leq i$. Then clearly for $\varphi_i^{-1}(g_i) = \left(\{g_i\} \times \prod_{j \in \omega, j \neq i} \{1, 2, \dots, m_j\} \right) \cap \varprojlim_{j \in \omega} G_j$

$$\psi^{-1}(\{l_0\} \times \{l_1\} \times \{l_2\} \times \dots \times \{l_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\}) = \varphi_i^{-1}(g_i),$$

which is open by the continuity of φ_i .

It is left to show that ψ is measure preserving. For the basic open set

$$B = \{l_0\} \times \{l_1\} \times \{l_2\} \times \dots \times \{l_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\} \in \mathcal{B},$$

let $g_i \in G_i$ be such that

$$\psi^{-1}(B) = \left(\{g_i\} \times \prod_{j \in \omega, j \neq i} \{1, 2, \dots, m_j\} \right) \cap \varprojlim_{j \in \omega} G_j = \varphi_i^{-1}(g_i).$$

Now, observe that if $g_i, h_i \in G_i$, and $h \in G$ is such that $\varphi_i(h) = h_i$, then $h\varphi_i^{-1}(g_i) = \varphi_i^{-1}(g_i h_i)$, i.e. sets of the form $\varphi_i^{-1}(x)$ ($x \in G_i$) are translates of each other. Using this, by the left-invariance of μ_G

$$\mu_G(\varphi_i^{-1}(g_i)) = \frac{1}{|G_i|} = \frac{1}{m_0} \cdot \frac{1}{m_1} \cdot \dots \cdot \frac{1}{m_i} = \nu(B).$$

Thus, we got that ν and $\mu \circ \psi^{-1}$ coincide on sets of the form $\{k_0\} \times \{k_1\} \times \dots \times \{k_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\}$ (i.e. on the elements of $\mathcal{B}$), from which they coincide on the

generated ring, since every finite union can be written as a disjoint union. Now, from [10, Sec. 13, Thm A], ν and $\mu \circ \psi^{-1}$ coincide on the generated σ -ring, that is the σ -algebra of the Borel sets, since the whole product $\prod_{j \in \omega} \{1, 2, \dots, m_j\}$ was contained in the ring. $\square$

For an arbitrary set X , let $\mathcal{P}(X)$ denote the power set of X . Now fix an $i \in \omega$. We define the following operations:

$$\begin{aligned}\mathcal{F}_i &: \mathcal{P}(G_i) \rightarrow \mathcal{P}(G_i) \\ \mathcal{F}_i &: H \mapsto HH \cup (HH)^{-1} \\ \mathcal{G}_i &: \mathcal{P}(G_i) \rightarrow \mathcal{P}(G_i) \\ \mathcal{G}_i &: H = \{g_1^{(j_1)}, g_2^{(j_2)}, \dots, g_n^{(j_n)}\} \mapsto \{g^{(j_k)} : g \in G_{i-1}, k \leq n\}.\end{aligned}$$

So $\mathcal{G}_i(H)$ is the minimal cover of H by the union of sets of the form $G_i^{(j)}$.

It is obvious that for $H \subseteq G_i$ we have

$$|\mathcal{F}_i(H)| \leq 2|H|^2$$

and

$$|\mathcal{G}_i(H)| \leq |H||G_{i-1}|,$$

thus

$$|\mathcal{G}_i \circ \mathcal{F}_i(H)| \leq 2|H|^2|G_{i-1}|.$$

From these inequalities it can be easily seen that for each $H \subseteq G_i$ there is an upper bound for $|(\mathcal{G}_i \circ \mathcal{F}_i)^j(H)|$ depending only on $|H|$, $|G_{i-1}|$, and j .

Now we define sets $B_i^j \subseteq \{1, 2, \dots, m_i\}$, for all pairs i, j such that

$$\text{if } k, k' \in B_i^j \Rightarrow G_i^{(k)} \cdot G_i^{(k')} \subseteq \{G_i^{(l)} : l \in B_i^{j+1}\} \quad (18)$$

and

$$\text{if } k \in B_i^j \Rightarrow \left(G_i^{(k)}\right)^{-1} \subseteq \{G_i^{(l)} : l \in B_i^{j+1}\} \quad (19)$$

i.e.

$$\{g^{(k)}(g')^{(k')} : k, k' \in B_i^j, g, g' \in G_{i-1}\} \subseteq \{g^{(l)} : l \in B_i^{j+1}, g \in G_{i-1}\}$$

and

$$\{(g^{(k)})^{-1} : k \in B_i^j, g \in G_{i-1}\} \subseteq \{g^{(l)} : l \in B_i^{j+1}, g \in G_{i-1}\}.$$

Let $A_i^0 = \mathcal{G}_i(\{e_{G_i}\}) \subseteq G_i$, and define $A_i^j = (\mathcal{G}_i \circ \mathcal{F}_i)^j(A_i^0)$ for each $j \leq i$, and let $A_i^j = G_i$, $B_i^j = \{1, 2, \dots, m_i\}$ for each $j > i$. For every j , as A_i^j is the union of sets of the form $G_i^{(k)}$, let

$$B_i^j = \{k : k \leq m_i, G_i^{(k)} \subseteq A_i^j\} \subseteq \{1, 2, \dots, m_i\} \quad (20)$$

(B_j^i stores the information: which elements of the partition $G_i = G_i^{(1)} \cup G_i^{(2)} \cup \dots \cup G_i^{(m_i)}$ are contained in A_i^j). By the construction of the A_i^j -s,

$$A_i^j A_i^j \subseteq A_i^{j+1}, \quad (21)$$

and

$$(A_i^j)^{-1} \subseteq A_i^{j+1}. \quad (22)$$

And $e_{G_i} \in A_i^0$ implies that for each $j \in \omega$, $e_{G_i} \in A_i^j$, hence

$$A_i^j \subseteq A_i^{j+1} \quad (j \in \omega) \quad (23)$$

Thus it is straightforward to check that (20) implies that B_i^j -s satisfy (18) and (19), and

$$B_i^j \subseteq B_i^{j+1} \quad (j \in \omega). \quad (24)$$

Next, we prove that by thinning out the sequence of the groups, we can assume that

$$|B_i^i| i^2 \leq m_i. \quad (25)$$

If the ω -type inverse system of finite groups $H_0 = \{e\}$, $H_1, H_2, \dots$ is given (where each H_i is a surjective homomorphic image of H_{i+1}), then one can construct a subsequence $(G_i = H_{n_i})_{i \in \omega}$ (where $n_0 = 0$), such that $|B_i^i| i^2 \leq m_i$ holds with the B_i^j -s and m_i -s defined for the system $(G_i)_{i \in \omega}$. The recursion goes as follows. Let $n_0 = 0$, and if $n_0, n_1, \dots, n_{i-1}$ are defined, then recall that for any $j \leq i$ there is an upper bound

$$d \geq |A_i^j| = |(\mathcal{G}_i \circ \mathcal{F}_i)^j(\{e\})|, \quad (26)$$

where d depends only on $|G_{i-1}| = |H_{n_{i-1}}|$ and j , no matter what G_i is. Thus we can choose n_i so that

$$di^2 \leq \frac{|H_{n_i}|}{|H_{n_{i-1}}|} \quad (27)$$

holds. Then, if $G_i = H_{n_i}$, using (27), (26) and the equality $|A_i^i| = |B_i^i| |G_{i-1}|$

$$m_i = \frac{|G_i|}{|G_{i-1}|} = \frac{|H_{n_i}|}{|H_{n_{i-1}}|} \geq di^2 \geq |A_i^i| i^2 \geq |B_i^i| i^2$$

thus (25) holds, indeed.

Now fix a non-principal ultrafilter $\mathcal{U}$ on ω . Define the subset $S' \subseteq \prod_{i \in \omega} \{1, 2, \dots, m_i\}$ as follows. For each $t \in \prod_{i \in \omega} \{1, 2, \dots, m_i\}$

$$t \in S' \iff \exists n : \{i : t_i \in B_i^n\} \in \mathcal{U}.$$

Let $S = \psi^{-1}(S')$. We have to check that

(i) S is a subgroup.

For $g = (g_i)_{i \in \omega} \in S$, let $s = \psi(g) \in \prod_{i \in \omega} \{1, 2, \dots, m_i\}$ (which means that

$$g = (g_i)_{i \in \omega} \quad g_i^{s_{i+1}} = g_{i+1}, \quad \forall i \in \omega$$

by (16), where we defined the ψ_i -s). Then obviously

$$g^{-1} = (g_i^{-1})_{i \in \omega},$$

and let $s' = \psi(g^{-1})$, thus again from the definition of ψ_i -s

$$(g_{i-1}^{-1})^{s'_i} = g_i^{-1}. \quad (28)$$

Choose a positive integer n_s such that $U_s = \{i : s_i \in B_{n_s}^i\} \in \mathcal{U}$. Now, for $i \in \omega$

$$i \in U_s \iff s_i \in B_{n_s}^i$$

which is equivalent to (using $g_{i-1}^{s_i} = g_i$ and (20))

$$s_i \in B_{n_s}^i \iff g_i \in A_{n_s}^i$$

and the latter combined with (21) implies

$$g_i \in A_{n_s}^i \Rightarrow g_i^{-1} \in A_{n_s+1}^i.$$

Now using (28) and (20) we get

$$g_i^{-1} \in A_{n_s+1}^i \iff s'_i \in B_{n_s+1}^i,$$

which means that $n_s + 1$ and U_s witness that $s' \in S'$, i.e. $\psi^{-1}(s') = g^{-1} \in S$.

In order to show that S is closed under multiplication, let $h \in S$ be another element, and $t = \psi(h)$ (i.e. $h = (h_i)_{i \in \omega}$, where $h_i \in G_i$, $h_i^{t_{i+1}} = h_{i+1}$), let n_t denote a positive integer for which $U_t = \{i : t_i \in B_{n_t}^i\} \in \mathcal{U}$. and let $n = \max\{n_s, n_t\}$, $U = U_s \cap U_t$. Suppose that $gh = (g_i h_i)_{i \in \omega}$ is represented by the sequence of integers $\psi(gh) = (v_i)_{i \in \omega}$. Then by (23)

$$A_{n_s}, A_{n_t} \subseteq A_n^i,$$

thus for any $i \in U$

$$g_i, h_i \in A_n^i.$$

Using (21)

$$g_i, h_i \in A_n^i \Rightarrow g_i h_i \in A_{n+1}^i,$$

and by $\psi(gh) = (v_i)_{i \in \omega}$ and (20),

$$g_i h_i \in A_{n+1}^i \iff v_i \in B_{n+1}^i,$$

thus $n + 1$ and U witness that $gh \in S$.

(ii) S is null.

Since the fixed bijection ψ between G and $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$ is a measure preserving homeomorphism, we can work in this product space, and it is enough to prove that S' is null.

$$S' \subseteq \bigcup_{n \in \omega} \bigcap_{j \in \omega} \bigcup_{k \geq j} \{s : s_k \in B_n^k\} = \bigcup_{n \in \omega} \{s : \{k : s_k \in B_n^k\} \text{ is infinite}\}$$

By σ -additivity, showing that for arbitrary fixed n the set $\bigcap_{j \in \omega} \bigcup_{k \geq j} \{s : s_k \in B_n^k\}$ is null will suffice.

Recall that if $l \leq l'$, then (by (24)) $B_k^l \subseteq B_k^{l'}$, and by (25) $|B_k^k| k^2 \leq m_k$, hence $\nu(\{s : s_k \in B_k^k\}) = \frac{|B_k^k|}{m_k} \leq \frac{1}{k^2}$. So if $k \geq n$, then

$$\nu(\{s : s_k \in B_k^n\}) \leq \frac{1}{k^2}.$$

Then, for arbitrary n, i if $i \geq n$,

$$\nu \left(\bigcap_{j \in \omega} \bigcup_{k \geq j} \{s : s_k \in B_k^n\} \right) \leq \nu \left(\bigcup_{k \geq i} \{s : s_k \in B_k^n\} \right) \leq \sum_{k=i}^{\infty} \frac{1}{k^2}.$$

Thus letting i tend to infinity we obtain

$$\nu \left(\bigcap_{j \in \omega} \bigcup_{k \geq j} \{s : s_k \in B_k^n\} \right) = 0.$$

(iii) S is not meager.

We work in the product space $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$ again, i.e. we prove that S' is of second category. Let R be a co-meager set in $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$, we will prove that $S' \cap R$ is nonempty.

Since $X = \prod_{i \in \omega} \{1, 2, \dots, m_i\}$ is a Polish space, thus there is a compatible complete metric d . The sets of the form

$$C_{p_0 p_1 p_2 \dots p_i} = \{p_0\} \times \{p_1\} \times \dots \times \{p_i\} \times \prod_{j \in \omega, j > i} \{1, 2, \dots, m_j\}$$

are compact open sets, and for each infinite sequence $(p_i)_{i \in \omega}$

$$\bigcap_{j \in \omega} C_{p_0 p_1 p_2 \dots p_j} = \{(p_i)_{i \in \omega}\},$$

therefore we can apply Lemma 24. Hence there is an increasing sequence $0 = n_0 < n_1 < n_2 < \dots$, and an $r \in \prod_{i \in \omega} \{1, 2, \dots, m_i\}$ such that for any $s \in \prod_{i \in \omega} \{1, 2, \dots, m_i\}$ if

$$\{j : s|_{[n_j, n_{j+1})} = r|_{[n_j, n_{j+1})}\} \text{ is infinite,}$$

then

$$\bigcap_{i \in \omega} C_{s_0 s_1 \dots s_i} = \bigcap_{i \in \omega} \left(\{s_0\} \times \{s_1\} \times \dots \times \{s_i\} \times \prod_{j > i} \{1, 2, \dots, m_j\} \right) \subseteq R$$

i.e. $s = (s_i)_{i \in \omega} \in R$.

Now one of the following sets:

$$\bigcup_{j \in \omega} [n_{2j}, n_{2j+1}), \quad \bigcup_{j \in \omega} [n_{2j+1}, n_{2j+2})$$

is in $\mathcal{U}$, let U denote that set. Now the following sequence

$$s_i = \begin{cases} 1 & \text{if } i \in U \\ r_i & \text{if } i \notin U \end{cases}$$

is obviously in R , because there are infinitely many intervals $[n_j, n_{j+1})$ contained in $\omega \setminus U$. On the other hand, by (17) we know that for the identity

$$e = (e_{G_i})_{i \in \omega} \in G$$

$\psi(e)$ is the constant 1 sequence in $\prod_{i \in \omega} \{1, 2, \dots, m_i\}$. Therefore $B_i^0 = \{1\}$ for all i , thus

$$\{i : s_i \in B_i^0\} \supseteq U \in \mathcal{U},$$

from which $s \in S$. We have that S is not disjoint from R . $\square$

Now we can turn to the case of connected Lie groups.

Proposition 27. *Let G be a second countable Lie group of positive dimension. Then there is a subgroup $S \leq G$ that is null, but non-meager.*

Proof. If G is d -dimensional, then let $C = [0, 1]^d = I^d$ denote the d dimensional unit cube, and let $B \supseteq I^d$ denote a ball containing it. Second countable Lie groups are Lindelöf spaces, thus there are open sets U_i ($1 \leq i \in \omega$) homeomorphic to B (by homeomorphisms $\psi_i : U_i \rightarrow B$), such that $\cup_{1 \leq i \in \omega} \psi_i^{-1}(\text{int}(I^d))$ covers G . Then also the $\psi_i^{-1}(I^d)$ -s (which may be overlapping) cover the group. We can assume that $e \in \psi_1^{-1}(I^d)$. Let φ_i be the restriction of ψ_i to the preimages of the cube (i.e. $\text{ran}(\varphi_i) = I^d$)

$$\begin{aligned} \varphi_i : \psi_i^{-1}(I^d) &\rightarrow I^d && \text{homeomorphism,} \\ x &\mapsto \psi_i(x). \end{aligned}$$

Let $\mathfrak{C}^i = \{C_1^i, C_2^i, \dots, C_{2^i}^i\}$ denote the i -th generation dyadic subdivision of the cube C , i.e. each C_j^i is of the form $[\frac{k_1}{2^i}, \frac{k_1+1}{2^i}] \times [\frac{k_2}{2^i}, \frac{k_2+1}{2^i}] \times \dots \times [\frac{k_d}{2^i}, \frac{k_d+1}{2^i}]$, where each $k_l \in \{0, 1, \dots, 2^i - 1\}$. From now on, if a set $X \subseteq G$ is given, $\varphi_k[X]$ denotes the image of X under φ_k , i.e.

$$\varphi_k[X] = \{\varphi_k(x) : x \in X \cap \text{dom}(\varphi_k)\}.$$

We will define an increasing sequence $(m_i)_{i \in \omega}$, and for each i an increasing sequence of subsets of $\mathfrak{C}^{m_i}$

$$D_0^i \subseteq D_1^i \subseteq \dots \subseteq D_j^i \subseteq \dots$$

such that for every $i \in \omega$, $i \geq 1$

- (i) $\cup D_j^i \subseteq \cup D_{j+1}^i$ for each $j \in \omega$,
- (ii) $e \in \varphi_1^{-1}(\cup D_0^i)$,
- (iii) for each $C^{m_{i-1}} \in \mathfrak{C}^{m_{i-1}}$ there exists $C^{m_i} \in D_0^i$ with $C^{m_i} \subseteq C^{m_{i-1}}$,
- (iv) for each $l \leq i$ and $j \in \omega$

$$\varphi_l \left[\left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup D_j^i) \right) \cdot \left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup D_j^i) \right) \right] \subseteq \cup D_{j+1}^i,$$

in other words, if $x \in \varphi_k^{-1}(\cup D_j^i)$, $y \in \varphi_{k'}^{-1}(\cup D_j^i)$ ($k, k' \leq i$), and $l \leq i$ is an integer for which $xy \in \text{dom}(\varphi_l)$, then $\varphi_l(xy) \in \cup D_{j+1}^i$,

- (v) for each $l \leq i$, $j \in \omega$ the following inclusion relation holds

$$\varphi_l \left[\left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup D_j^i) \right)^{-1} \right] \subseteq \cup D_{j+1}^i$$

in other words, if $k \leq i$, where $x \in \varphi_k^{-1}(\cup D_j^i)$, and $l \leq i$ is an integer for which $x^{-1} \in \text{dom}(\varphi_l)$, then $\varphi_l(x^{-1}) \in \cup D_{j+1}^i$,

- (vi) for every $r \leq i$

$$\mu \left(\varphi_r^{-1} \left(\bigcup D_j^i \right) \right) \leq \frac{1}{i^2}. \quad (29)$$

Lemma 28. *There are such $(m_i)_{i \in \omega, i \geq 1}$, $(D_j^i)_{1 \leq i \in \omega, j \in \omega}$ sequences.*

Proof. Let $m_0 = 0$, and $D_j^0 = C = I^d$ for every $j \in \omega$.

Assume that $m_1 < m_2 < \dots < m_{i-1}$, and $D_k^j \subseteq \mathfrak{C}^{m_k}$ ($j < i, k \in \omega$) are defined.

First we define the D_j^i -s if $j \leq i$. Let $x_0 = \varphi_1(e) \in C^{m_{i-1}} \in \mathfrak{C}^{m_{i-1}}$, and define the finite set $A_0 \subseteq I^d$ by translating x_0 to each $C^{m_{i-1}} \in \mathfrak{C}^{m_{i-1}}$

$$A_0 = \left\{ x_0 + \left(\frac{s}{2^{m_{i-1}}}, \frac{t}{2^{m_{i-1}}} \right) : -2^{m_{i-1}} \leq s, t \leq 2^{m_{i-1}} \right\} \cap I^d.$$

Define $B_0 = \bigcup_{j=1}^i \varphi_j^{-1}(A_0) \subseteq G$

Define the following operation $\mathcal{F}$ on $\mathcal{P}(G)$

$$\begin{aligned} \mathcal{F} : \mathcal{P}(G) &\rightarrow \mathcal{P}(G) \\ H &\mapsto (H \cup H^{-1})^2 \end{aligned} \quad (30)$$

Define inductively the sequences B_j, A_j ($j \leq i$) by the following procedure. If B_{j-1}, A_{j-1} are given, then

$$A_j = \bigcup_{k=1}^i \varphi_k[\mathcal{F}(B_{j-1})] \quad (31)$$

i.e. we choose $A_j \subseteq I^d$ by taking for each element of $\mathcal{F}(B_{j-1})$ all corresponding points in the a cube according to the φ_k -s ($k \leq i$). (Notice that there may be points in $\mathcal{F}(B_{j-1})$ which are not contained in $\text{dom}(\varphi_k)$ -s ($k \leq i$), thus there are no corresponding points.) And B_j will be the set of corresponding group elements according to the first i charts

$$B_j = \bigcup_{k=1}^i \varphi_k^{-1}(A_j). \quad (32)$$

By induction on j , we can prove that $e \in B_j$. We know that $e \in B_0$, and if $e \in B_j$, then by (30) $e \in \mathcal{F}(B_j)$, thus using (31) and $i \geq 1$ $\varphi_1(e) \in A_{j+1}$, from which

$$e \in \bigcup_{k=1}^i \varphi_k^{-1}(A_{j+1}) = B_{j+1}. \quad (33)$$

Now (33) implies that

$$B_j \subseteq \mathcal{F}(B_j), \quad (34)$$

and it can be easily seen by induction on j that the B_j -s and the A_j -s form an increasing sequence. Indeed, for each $j \in \omega$ using (31), (34) and (32)

$$A_{j+1} = \bigcup_{k=1}^i \varphi_k[\mathcal{F}(B_j)] \supseteq \bigcup_{k=1}^i \varphi_k[B_j] = \bigcup_{k=1}^i \left(\varphi_k \left[\bigcup_{k=1}^i \varphi_k^{-1}(A_j) \right] \right) \supseteq A_j,$$

and from that

$$B_{j+1} = \bigcup_{k=1}^i \varphi_k^{-1}(A_{j+1}) \supseteq \bigcup_{k=1}^i \varphi_k^{-1}(A_j) = B_j.$$

For each $k \geq m_{i-1}$, one can cover the finite set A_i uniquely by a minimal, at most $2^d |A_i|$ -many element subset $E \subseteq \mathfrak{C}^k$, such that $A_i \subseteq \text{int}_{I^d}(\cup E^k)$ (where int_{I^d} denotes the interior with respect to the subspace topology of the cube). For each

$k \geq m_{i-1}$ let E^k denote that set. Clearly $\bigcup E^k \supseteq \bigcup E^{k+1}$, and $\bigcap_{k=m_{i-1}}^{\infty} \bigcup E^k = A_i$. This implies that, since φ_r -s are homeomorphisms

$$\bigcap_{k=m_{i-1}}^{\infty} \varphi_r^{-1}(\bigcup E^k) = \varphi_r^{-1}(A_i)$$

Which intersection is of measure zero, since finite, and G is of positive dimension. Using the Borel-Cantelli Lemma, for any $r \leq i$ there exists a k , such that

$$\mu\left(\varphi_r^{-1}\left(\bigcup E^k\right)\right) \leq \frac{1}{i^2}$$

Hence there exists a sufficiently large l_0 , which is appropriate for each $r \leq i$. Let $E_i = E^{l_0} \subseteq \mathfrak{C}^{l_0}$. Now beginning from $j = i$, and stepping down with j to 0, we will define a sequence $E_j \subseteq \mathfrak{C}^{l_j}$ for some l_j depending only on j and the already defined E_p -s ($p > j$, $p \leq i$), for which the following holds

- $\bigcup E_{j+1} \supseteq \bigcup E_j$,
- A_j is in the relative interior (i.e. in the cube) of $\bigcup E_j$:

$$\text{int}_{I^d}(\bigcup E_j) \supseteq A_j, \quad (35)$$

- if (for $r, s \leq i$) $x \in \varphi_r^{-1}(\bigcup E_j)$, $y \in \varphi_s^{-1}(\bigcup E_j)$ and $k \leq i$ is such that $xy \in \text{dom}(\varphi_k)$, then $\varphi_k(xy) \in \bigcup E_{j+1}$, in other words

$$(\forall r, s \leq i, k \leq i : \varphi_k[\varphi_r^{-1}(\bigcup E_j) \cdot \varphi_s^{-1}(\bigcup E_j)]) \subseteq \bigcup E_{j+1}. \quad (36)$$

- If $x \in \varphi_r^{-1}(\bigcup E_j)$ ($r \leq i$), and $t \leq i$ is such that $x^{-1} \in \text{dom}(\varphi_t)$ then $\varphi_t(x^{-1}) \in \bigcup E_{j+1}$, i.e.:

$$(\forall r \leq i, t \leq i : \varphi_t[(\varphi_r^{-1}(\bigcup E_j))^{-1}] \subseteq \bigcup E_{j+1} \quad (37)$$

Suppose that $E_{j+1}, E_{j+2}, \dots, E_i$ are already defined. Again, for each $l > m_{i-1}$ there is a unique minimal covering $E_j^l \subseteq \mathfrak{C}^l$ of A_j , $|E_j^l| \leq 2^d |A_j|$, for which (35) holds. Recall (32), i.e.

$$B_j = \bigcup_{k=1}^i \varphi_k^{-1}(\bigcup A_j) .$$

Now, for each $x, y \in A_j$ we would like to choose a sufficiently small neighborhoods U_x, V_y in the cube so that whenever $r, s \leq i$, choosing $z = \varphi_r^{-1}(x)$, $w = \varphi_s^{-1}(y)$, and $B_z = \varphi_r^{-1}(U_x)$, $B_w = \varphi_s^{-1}(V_y)$, then for any $t \leq i$

$$\varphi_t[B_z B_w] \subseteq \bigcup E_{j+1}$$

i.e.

$$p \in B_z B_w \cap \text{dom}(\varphi_t) \Rightarrow \varphi_t(p) \in \bigcup E_{j+1}.$$

Equivalently, for each $x, y \in A_j$

$$\varphi_t[\varphi_r^{-1}(U_x) \varphi_s^{-1}(V_y)] \subseteq \bigcup E_{j+1} \quad (\text{for each } r, s, t \leq i). \quad (38)$$

First for any fixed $r, s, t \leq i$ we choose open neighborhoods $U_{r,x,s,y}^t$ of x , and $V_{r,x,s,y}^t$ of y such that

$$\varphi_t[\varphi_r^{-1}(U_{r,x,s,y}^t) \varphi_s^{-1}(V_{r,x,s,y}^t)] \subseteq \bigcup E_{j+1} \quad (39)$$

holds. Let $x, y \in A_j$, $r, s, t \leq i$ be fixed. Then, if $z = \varphi_r^{-1}(x)$, $w = \varphi_s^{-1}(y)$,

- if $t \leq i$ is such that $zw \in \text{dom}(\varphi_t)$,
then, since $z, w \in B_j$, $zw \in \mathcal{F}(B_j)$ (by (30)), hence $a = \varphi_t(zw) \in A_{j+1}$ (by (31)). By the construction of E^{j+1} , $a \in \text{int}_{I^d}(\cup E_{j+1})$ (recall that E_{j+1} was chosen so that $A_{j+1} \subseteq \text{int}_{I^d}(\cup E_{j+1})$). Since φ_t is a homeomorphism, $\varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1}))$ is relatively open in $\text{dom}(\varphi_t)$. Therefore there is an open set W in G containing zw , such that

$$W \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1})). \quad (40)$$

Then, by the continuity of the multiplication, we have that there are open sets $U, V \subseteq G$, $z \in U$, $w \in V$, for which

$$UV \subseteq W. \quad (41)$$

Now we claim that

$$\varphi_t[(U \cap \text{dom}(\varphi_r)) \cdot (V \cap \text{dom}(\varphi_s))] \subseteq \text{int}_{I^d}(\cup E_{j+1}). \quad (42)$$

Indeed, let $g \in (U \cap \text{dom}(\varphi_r)) \cdot (V \cap \text{dom}(\varphi_s))$, $g \in \text{dom}(\varphi_t)$ be fixed. Then by (40) and (41)

$$g \in W \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1})),$$

thus

$$\varphi_t(g) \in \text{int}_{I^d}(\cup E_{j+1}),$$

therefore (42) holds.

Let $U_{r,x,s,y}^t = \varphi_r[U] \ni x$, $V_{r,x,s,y}^t = \varphi_s[V] = \varphi_s[V \cap \text{dom}(\varphi_s)] \ni y$ relatively open sets in the cube, then (42) implies that

$$\varphi_t[\varphi_r^{-1}(U_{r,x,s,y}^t)\varphi_s^{-1}(V_{r,x,s,y}^t)] \subseteq \text{int}_{I^d}(\cup E_{j+1}).$$

- if t is such that $zw \notin \varphi_t^{-1}(I^d)$,
then (recall that φ_t is a homeomorphism, thus $\varphi_t^{-1}(I^d)$ is compact), there are open sets $U \ni z$, $V \ni w$ in G such that

$$UV \cap \varphi_t^{-1}(I^d) = \emptyset. \quad (43)$$

Let $U_{r,x,s,y}^t = \varphi_r[U] \ni x$, $V_{r,x,s,y}^t = \varphi_s[V] \ni y$ be open sets in the cube, then (43) clearly yields that

$$\varphi_r^{-1}(U_{r,x,s,y}^t)\varphi_s^{-1}(V_{r,x,s,y}^t \cap \text{dom}(\varphi_t)) = \emptyset,$$

thus

$$\varphi_t[\varphi_r^{-1}(U_{r,x,s,y}^t)\varphi_s^{-1}(V_{r,x,s,y}^t)] = \emptyset.$$

In both cases (39) holds.

Now we can define the U_x -s ($x \in A_j$) and V_y -s ($y \in A_j$) as

$$U_x = \left(\bigcap_{s,r,t \leq i, y \in A_j} U_{r,x,s,y}^t \right), \quad (44)$$

and

$$V_y = \bigcap_{s,r,t \leq i, x \in A_j} V_{r,x,s,y}^t. \quad (45)$$

Then $U_x \ni x$, $V_y \ni y$ are relatively open sets in the cube (since A_j is finite), and clearly for arbitrary $r, s, t \leq i$, by (44), (45) and (39)

$$\begin{aligned} \varphi_r^{-1}(U_x) \cdot \varphi_s^{-1}(V_y) \cap \text{dom}(\varphi_t) &\subseteq (\varphi_r^{-1}(U_{r,x,s,y}^t) \cdot \varphi_s^{-1}(V_{r,x,s,y}^t)) \cap \text{dom}(\varphi_t) \subseteq \\ &\subseteq \varphi_t^{-1}(\cup E_{j+1}). \end{aligned}$$

Now we would like to find an open neighborhood $W_x \subseteq I^d$ of every $x \in A_j$ such that for each $r, t \leq i$

$$(\varphi_r^{-1}(W_x))^{-1} \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\cup E_{j+1}). \quad (46)$$

First for each $x \in A_j$ and $r, t \leq i$ we define neighborhoods $W_{x,r}^t$ of x in I^d such that

$$(\varphi_r^{-1}(W_{x,r}^t))^{-1} \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\cup E_{j+1}). \quad (47)$$

Let $x \in A_j$ and $r, t \leq i$ be fixed. Set $z = \varphi_r^{-1}(x)$.

- If $z^{-1} = (\varphi_r^{-1}(x))^{-1} \in \text{dom}(\varphi_t)$, then
since $z \in B_j$, $z^{-1} \in \mathcal{F}(B_j)$ (by (30)), and $z^{-1} \in \text{dom}(\varphi_t)$ by (31), $\varphi_t(z^{-1}) \in A_{j+1}$, thus by (32) $z^{-1} \in B_{j+1}$.
We know that $A_{j+1} \subseteq \text{int}_{I^d}(\cup E_{j+1})$, i.e. z^{-1} is an element of the relatively open set $\varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1})) \subseteq \text{dom}(\varphi_t)$, from which there is an open $V \subseteq G$ containing z^{-1} such that

$$V \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1})).$$

Now, there is an open neighborhood $W' \subseteq G$ of z such that $(W')^{-1} \subseteq V$. If we set $W = \varphi_r[W'] = \varphi_r[W' \cap \text{dom}(\varphi_r)]$, then $x \in W$ and

$$(\varphi_r^{-1}(W))^{-1} \subseteq (W')^{-1} \subseteq V$$

from which it follows that

$$(\varphi_r^{-1}(W))^{-1} \cap \text{dom}(\varphi_t) \subseteq V \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\text{int}_{I^d}(\cup E_{j+1})),$$

i.e.

$$\varphi_t[(\varphi_r^{-1}(W))^{-1}] \subseteq \text{int}_{I^d}(\cup E_{j+1}).$$

- If $z^{-1} = (\varphi_r^{-1}(x))^{-1} \notin \text{dom}(\varphi_t)$, then by the compactness of $\text{dom}(\varphi_t)$, there is an open neighborhood V of z^{-1} such that $V \cap \text{dom}(\varphi_t) = \emptyset$. But then choosing $W' = V^{-1}$, W' is a neighborhood of z such that $(W')^{-1} = V \subseteq G \setminus \text{dom}(\varphi_t)$. Then, with $W = \varphi_r[W'] = \varphi_r[W' \cap \text{dom}(\varphi_r)]$ clearly $x \in W$ and

$$(\varphi_r^{-1}(W))^{-1} \subseteq (W')^{-1} \subseteq V,$$

thus $(\varphi_r^{-1}(W))^{-1} \cap \text{dom}(\varphi_t) = \emptyset$.

In both cases i.e. independently of whether $z^{-1} \in \text{dom}(\varphi_t)$, let $W_{x,r}^t = W$.

Now we can define the W_x -s ($x \in A_j$)

$$W_x = \bigcap_{r,t \leq i} W_{x,r}^t. \quad (48)$$

Then, fixing $x \in A_j$, for any $r, t \leq i$, using (48) and (47)

$$(\varphi_r^{-1}(W_x))^{-1} \cap \text{dom}(\varphi_t) \subseteq (\varphi_r^{-1}(W_{x,r}^t))^{-1} \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\cup E_{j+1}),$$

i.e. (46) holds, indeed.

Now we have that A_j is contained in the open sets $\bigcup_{x \in A_j} U_x$, $\bigcup_{y \in A_j} V_y$ and $\bigcup_{x \in A_j} W_x$.

For each $p \geq m_{i-1}$, let $E^p \subseteq \mathfrak{C}^p$ denote the unique minimal covering of A_j , for which $A_j \subseteq \text{int}_{I^d}(\cup E^p)$, and hence $|E^k| \leq 2^d |A_j|$. Therefore, if p is sufficiently large, then

$$A_j \subseteq \cup E^p \subseteq \left(\bigcup_{x \in A_j} U_x \right) \cap \left(\bigcup_{y \in A_j} V_y \right) \cap \left(\bigcup_{x \in A_j} W_x \right) \quad (49)$$

Furthermore, we can choose p large enough so that

$$\cup E^p \subseteq \cup E_{j+1}, \quad (50)$$

since we saw that A_j -s form an increasing sequence, i.e. $A_j \subseteq A_{j+1}$. Fix such a p_0 for which (49) and (50) hold, and let $l_j = p_0$, $E_j = E^{l_j}$.

Now we check that (36) holds, i.e. for any $t \leq i$

$$\varphi_t \left[\left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup E^p) \right) \left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup E^p) \right) \right] \subseteq \cup E_{j+1}.$$

For any $r, s \leq i$, since $\cup E^p \subseteq \left(\bigcup_{x \in A_j} U_x \right)$, and $\cup E^p \subseteq \left(\bigcup_{y \in A_j} V_y \right)$

$$\varphi_t \left[\varphi_r^{-1}(\cup E^p) \varphi_s^{-1}(\cup E^p) \right] \subseteq \varphi_t \left[\varphi_r^{-1} \left(\bigcup_{x \in A_j} U_x \right) \varphi_s^{-1} \left(\bigcup_{y \in A_j} V_y \right) \right]$$

and using (38)

$$\varphi_t \left[\varphi_r^{-1} \left(\bigcup_{x \in A_j} U_x \right) \varphi_s^{-1} \left(\bigcup_{y \in A_j} V_y \right) \right] = \bigcup_{x,y \in A_j} \varphi_t \left[\varphi_r^{-1}(U_x) \varphi_s^{-1}(V_y) \right] \subseteq \cup E_{j+1}.$$

Similarly, since $\cup E^p \subseteq \left(\bigcup_{x \in A_j} W_x \right)$, for any fixed $t \leq i$

(37) holds, i.e.

$$\left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup E^p) \right)^{-1} \subseteq \left(\bigcup_{k=1}^i \varphi_k^{-1} \left(\bigcup_{x \in A_j} W_x \right) \right)^{-1} = \bigcup_{k=1}^i \bigcup_{x \in A_j} (\varphi_k^{-1}(W_x))^{-1},$$

and taking its intersection with $\text{dom}(\varphi_t)$, using (46)

$$\left(\bigcup_{k=1}^i \varphi_k^{-1}(\cup E^p) \right)^{-1} \cap \text{dom}(\varphi_t) \subseteq \bigcup_{k=1}^i \bigcup_{x \in A_j} (\varphi_k^{-1}(W_x))^{-1} \cap \text{dom}(\varphi_t) \subseteq \varphi_t^{-1}(\cup E_{j+1}).$$

At the point when E_j is already defined for all $0 \leq j \leq i$, and l_0 denotes the fineness of E_0 , i.e. $E_0 \subseteq \mathfrak{C}^{l_0}$, then let $m_i = l_0$, and for each $j \leq i$ let $D_j^i \subseteq \mathfrak{C}^{m_i}$ such that $\cup D_j^i = \cup E_j$, i.e. a subdivision of E_j . Since (36) and (37) hold, and because $\cup E_j = \cup D_j^i$ ($j \leq i$), (iv) and (v) obviously hold for the D_j^i -s where $j < i$. (2) implies (i).

For $j > i$, let $D_j^i = \mathfrak{C}^{m_i}$, i.e. we choose all the 2^{m_i} -many small cubes, thus (iv) and (v) hold in this case too. (i) is also true. Since $\text{int}_{I^d}(\cup E_0^i) \supseteq A_0$, and A_0 contains at least one point from each $C \in \mathfrak{C}^{m_{i-1}}$, $\text{int}_{I^d}(\cup D_0^i) = \text{int}_{I^d}(\cup E_0^i)$ intersects each $C \in \mathfrak{C}^{m_{i-1}}$. Thus for each $C \in \mathfrak{C}^{m_{i-1}}$, D_0^i contains a cube C' such that $C' \subseteq C$, i.e. (iii) holds. □

Now we are ready to construct the desired subgroup.

For each $z \in I^d$ let B_z denote the set of the following (corresponding) subset of $\prod_{i \in \omega} \mathfrak{C}^{m_i}$.

$$B_z = \{(C^{m_i})_{i \in \omega} \mid \cap C^{m_i} = \{z\}\}$$

Now choose a non-principal ultrafilter $\mathcal{U}$ on ω , and define $S' \subseteq I^d$ in the following way:

$$S' = \{z \mid \exists l \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_l^i\} \in \mathcal{U}\}$$

Let $S = \bigcup_{i=1}^\infty \varphi_i^{-1}(S')$ We will show that S is a subgroup of G , S is null, and is not meager:

(i) S is a subgroup.

- First, S is closed under the multiplication. Let $x, y \in S$
and $k_1, k_2, k_3 \in \omega$ such that $x \in \text{dom}(\varphi_{k_1})$, $y \in \text{dom}(\varphi_{k_2})$, $xy \in \text{dom}(\varphi_{k_3})$, $\varphi_{k_1}(x) = x'$, $\varphi_{k_2}(y) = y'$ and let $(E_x^{m_i})_{i \in \omega} \in B_{x'}$, $(E_y^{m_i})_{i \in \omega} \in B_{y'}$ $U_x, U_y \in \mathcal{U}$, l_x, l_y such that

$$\{i : E_x^{m_i} \in D_{l_x}^i\} = U_x \tag{51}$$

$$\{i : E_y^{m_i} \in D_{l_y}^i\} = U_y. \tag{52}$$

Recall that $\cup D_k^i \subseteq \cup D_{k+1}^i$ by the construction of the D_k^i -s, see (i). Let $l = \max\{l_x, l_y, k_1, k_2, k_3\}$ denote the largest one, then if $i \in U_x \cap U_y \in \mathcal{U}$, using (51), (52)

$$x' = \varphi_{k_1}(x) \in E_x^{m_i} \in D_{l_x}^i \subseteq D_l^i$$

and

$$y' = \varphi_{k_2}(y) \in E_y^{m_i} \in D_{l_y}^i \subseteq D_l^i,$$

i.e.

$$\{i : E_x^{m_i} \in D_l^i\} \supseteq U_x \supseteq U_x \cap U_y \in \mathcal{U}, \quad (53)$$

$$\{i : E_y^{m_i} \in D_l^i\} \supseteq U_y \supseteq U_x \cap U_y \in \mathcal{U}. \quad (54)$$

From which,

$$xy \in (\varphi_{k_1}^{-1}(\cup D_l^i) \cdot \varphi_{k_2}^{-1}(\cup D_l^i)) \cap \text{dom}(\varphi_{k_3}).$$

But if i is such that $l_x, l_y, k_3 \leq l \geq i$, we can use the property (iv) of the D_j^i -s, that implies $xy \in \varphi_{k_3}^{-1}(\cup D_{l+1}^i)$, i.e. there is an $(E^{m_i})' \in D_{l+1}^i$, for which

$$\varphi_{k_3}(xy) \in (E^{m_i})'.$$

This is true for all $i \in U_x \cap U_y \in \mathcal{U}$, $i \geq l$, thus there is a partial sequence $((E^{m_i})')_{l \geq i \in U_x \cap U_y}$ for which

$$\varphi_{k_3}(xy) \in (E^{m_i})' \text{ for each } i \in U_x \cap U_y, i \geq l.$$

Choosing $(E^{m_i})'$ -s for the remaining i -s ($\omega \setminus (U_x \cap U_y \cap [l, \infty))$) so that $\varphi_{k_3}(xy) \in (E^{m_i})'$ hold will yield an appropriate sequence $((E^{m_i})')_{i \in \omega}$ from $B_{\varphi_{k_3}(xy)}$, i.e.

$$\{\varphi_{k_3}(xy)\} = \bigcap_{i \in \omega} (E^{m_i})',$$

and

$$\{i \in \omega : (E^{m_i})' \in D_{l+1}^i\} \supseteq U_x \cap U_y \cap [l, \infty).$$

Hence $((E^{m_i})')_{i \in \omega}$, $l+1$, $U_x \cap U_y \cap [l, \infty)$ witness that $\varphi_{k_3}(xy) \in S'$, we can conclude that

$$xy \in S.$$

- Similarly, for $x \in S$ if $x^{-1} \in \text{dom}(\varphi_k)$, and $\varphi_{k_1}(x) = x' \in S'$ is witnessed by the aforementioned $((E_{x'}^{m_i})_{i \in \omega}, l_x, U_x)$, then let $l = \max\{l_x, k_1\}$. Now, if $i \in U_x$, $i \geq l$:

$$\varphi_{k_1}(x) = x' \in E_x^{m_i} \in D_{l_x}^i \subseteq D_l^i$$

Then $x^{-1} \in \text{dom}(\varphi_k)$, thus by (v), and the conditions ($k \leq l \leq i$)

$$\varphi_k(x^{-1}) \in \cup D_{l+1}^i$$

so there is an $(E^{m_i})' \in D_{l+1}^i$ for which $\varphi_k(x^{-1}) \in (E^{m_i})'$. Again, this is true for each $i \in U_x$, $i \geq l$, i.e. we have a sequence $((E^{m_i})')_{i \in \omega}$ from $B_{\varphi_k(x^{-1})}$, such that for all $i \in U_x$ if $i \geq l$, then $(E^{m_i})' \in D_{l+1}^i$. Since $U_x \cap [l, \infty) \in \mathcal{U}$, this verifies that $\varphi_k(x^{-1}) \in S'$, hence $x^{-1} \in S$.

(ii) S is null.

By the σ -additivity, it is enough to show that $\mu(\varphi_p^{-1}(S')) = 0$ for each $p \in \omega$. (And since

$$S' = \bigcup_{l \in \omega} \{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_l^i\} \in \mathcal{U}\}$$

it is sufficient to show that for each l

$$\mu(\varphi_p^{-1}(\{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_l^i\} \in \mathcal{U}\})) = 0.)$$

But

$$\begin{aligned} \{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_l^i\} \in \mathcal{U}\} &\subseteq \\ \{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_i^i\} \in \mathcal{U}\} &\subseteq \end{aligned}$$

since for fixed l , $D_l^i \subseteq D_i^i$ is true for all but finitely many $(l-1)$ -many i -s, and $\mathcal{U}$ is non-principal. Moreover the latter set is contained in a larger one, corresponding to sequences $(C^{m_i})_{i \in \omega} \in \prod_{i \in \omega} \mathfrak{C}^{m_i}$ for which there exist infinitely many i -s such that $C^{m_i} \in D_i^i$

$$\{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_i^i\} \in \mathcal{U}\} \subseteq$$

$$\bigcap_{k=1}^{\infty} \bigcup_{l=k}^{\infty} \{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : C^{m_l} \in D_l^l\} \subseteq$$

$$\bigcap_{k=1}^{\infty} \bigcup_{l=k}^{\infty} \cup D_l^l$$

Considering its preimage under the homeomorphism φ_p , then for any $r \in \omega$

$$\varphi_p^{-1} \left(\bigcap_{k=1}^{\infty} \bigcup_{l=k}^{\infty} \cup D_l^l \right) \subseteq \varphi_p^{-1} \left(\bigcup_{l=r}^{\infty} \cup D_l^l \right)$$

and applying (29) (i.e. $t \geq p$ implies that $\mu(\varphi_p^{-1}(\cup D_t^t)) \leq \frac{1}{t^2}$) for $r \geq p$,

$$\begin{aligned} \mu(\varphi_p^{-1}(\{z \in I^d \mid \exists (C^{m_i})_{i \in \omega} \in B_z : \{i \in \omega \mid C^{m_i} \in D_i^i\} \in \mathcal{U}\})) &\leq \\ &\leq \mu \left(\varphi_p^{-1} \left(\bigcap_{k=1}^{\infty} \bigcup_{l=k}^{\infty} \cup D_l^l \right) \right) \leq \mu \left(\varphi_p^{-1} \left(\bigcup_{l=r}^{\infty} \cup D_l^l \right) \right) \leq \sum_{l=r}^{\infty} \frac{1}{l^2} \end{aligned}$$

which upper bound tends to 0 as r tends to ∞ .

(iii) S is not meager.

Let $R \subseteq G$ be co-meager. First we show that if $R \supseteq \bigcap_{i \in \omega} U_i$, where U_i -s are dense open sets, $R \cap \text{dom}(\varphi_1)$ is co-meager in $\text{dom}(\varphi_1)$. Since for each i $U_i \cap \text{dom}(\varphi_1)$ is open in $\text{dom}(\varphi_1)$, and dense in $\text{dom}(\varphi_1)$ (because $U_i \cap \text{dom}(\varphi_1)$ is dense in $\text{int}(\text{dom}(\varphi_1))$, and the latter is dense in $\text{dom}(\varphi_1)$), therefore $\bigcap_{i \in \omega} (U_i \cap \text{dom}(\varphi_1)) \subseteq R \cap \text{dom}(\varphi_1)$ is comeager in $\text{dom}(\varphi_1)$.

Since φ_1 is a homeomorphism, the image $\varphi_1[R] \subseteq I^d$ of $R \cap \text{dom}(\varphi_1)$ is co-meager. Now we can apply Lemma 24 in I^d , for the co-meager set $\varphi_1[R]$, with the compact sets $\bigcup_{i \in \omega} \mathfrak{C}^{m_i}$ (for each $C^{m_i} \in \mathfrak{C}^{m_i}$ we can assign (inductively) a sequence of the form $p_0 p_1 \dots p_i$, (where $1 \leq p_k \leq 2^{m_k - m_{k-1}}$) such that the obvious inclusion relation holds, i.e. $\mathfrak{C}^{m_k} = \{C_{p_0 p_1 \dots p_k} : 1 \leq p_j \leq 2^{m_j - m_{j-1}} (\forall j \leq k)\}$, and $C_{p_0 p_1 \dots p_k} = \bigcup_{1 \leq j \leq 2^{m_{k+1} - m_k}} C_{p_0 p_1 \dots p_k j}$). The lemma gives an increasing sequence $(n_j)_{j \in \omega}$, and a sequence $(r_j)_{j \in \omega}$, ($1 \leq r_j \leq 2^{m_j - m_{j-1}}$), such that for any infinite sequence $(s_i)_{i \in \omega}$ ($1 \leq s_l \leq 2^{m_l - m_{l-1}}$)

$$\text{if the set } \{j : r_{[n_j, n_{j+1})} = s_{[n_j, n_{j+1})}\} \text{ is infinite} \Rightarrow \bigcap_{i \in \omega} C_{s_0 s_1 \dots s_i} \subseteq \varphi_1[R].$$

Clearly, exactly one of the following sets is in $\mathcal{U}$:

$$\bigcup_{j \in \omega} [n_{2j}, n_{2j+1}) \quad \text{and} \quad \bigcup_{j \in \omega} [n_{2j+1}, n_{2j+2})$$

Let U denote that element of $\mathcal{U}$, and define the sequence $(E^{m_i})_{i \in \omega}$ by induction on i : let $E^{m_0} = E^0 = I^d$, and define $E^{m_{i+1}}$ (recall that by (iii), for all $C^{m_k} \in \mathfrak{C}^{m_k}$ there is an element $C \in D_0^{k+1} \subseteq \mathfrak{C}^{m_{k+1}}$ for which $C \subseteq C^{m_k}$) as

$$E^{m_{i+1}} = \begin{cases} C \in D_0^{i+1}, C \subseteq E^{m_i} & \text{if } i+1 \in U \\ C_{p_0 p_1 \dots p_i r_{i+1}} & \text{if } i+1 \notin U \text{ and } E^{m_i} = C_{p_0 p_1 \dots p_i} \end{cases}$$

Clearly, if $\{z\} = \bigcap_{i \in \omega} E^{m_i}$, then $z \in S'$, because

$$\{i : E^{m_i} \in D_0^i\} \supseteq U \in \mathcal{U}.$$

But $z \in \varphi_1[R]$ also, since if $p = p_0 p_1 \dots p_l \dots \in \{1\} \times \prod_{l \in \omega, l \geq 1} \{1, 2, \dots, 2^{m_l - m_{l-1}}\}$ is such that

$$(E^{m_i})_{i \in \omega} = (C_{p_0 p_1 \dots p_i})_{i \in \omega},$$

then because $\omega \setminus U$ contains infinitely many intervals of the form $[n_l, n_{l+1})$,

$$\{j : p|_{[n_j, n_{j+1})} = r|_{[n_j, n_{j+1})}\} \text{ is infinite,}$$

thus

$$z \in \bigcap_{i \in \omega} E^{m_i} = \bigcap_{i \in \omega} C_{p_0 p_1 \dots p_i} \subseteq \varphi_1[R].$$

From this, $\varphi_1^{-1}(z) \in S \cap R$.

□

Now we are ready to state our first main theorem.

Theorem 29. *If G is a locally compact group that is non-discrete, and μ is a left Haar measure then there is a subgroup $S \leq G$ for which $\mu(H) = 0$, but S is not meager.*

Proof. Using Lemma 12, there is an open subgroup $H \leq G$, in which for each neighborhood $U \subseteq H$ of the identity, there is a compact normal subgroup $N_U \subseteq U$ of H such that H/N_U is a Lie group having finitely many connected components (fix this operation $U \mapsto N_U$). Thus, using Corollary 16, H is an inverse limit of Lie groups all of which have finitely many connected components. Now we define a compact normal subgroup $K \triangleleft H$ such that H/K is a second countable Lie group of positive dimension, or an inverse limit of a countable sequence of finite groups, where the size of the groups form a strictly increasing sequence, thus (by Proposition 25 and 27) there is a null, but non-meager subgroup in H/K . Now,

- if there is an

$$N \in \mathfrak{N}_H = \{N \triangleleft H : H/N \text{ is a Lie group}$$

with finitely many connected components\}

for which H/N is a Lie group of positive dimension, then we are done.

- otherwise, each H/N ($N \in \mathfrak{N}_H$) is a finite group. Recall that, using Lemma 14 $\mathfrak{N}_H$ is closed under finite intersection. First, then there cannot exist a least $N_0 \in \mathfrak{N}_H$ (wrt. the inclusion), because the mapping $g \mapsto (gN)_{N \in \mathfrak{N}_H}$ (which maps H to $\varprojlim_{N \in \mathfrak{N}_H} H/N$) is an isomorphism by Corollary 16, thus its kernel is $\{e\} = \bigcap_{N \in \mathfrak{N}_H} N = N_0$ (by Lemma 15). This would imply that $H = H/N_0$ is a finite discrete group. But H was an open subgroup of G , thus G is discrete, which is a contradiction.

Then there cannot exist a minimal $N \in \mathfrak{N}_H$ either, because that would be a least element, since $\mathfrak{N}_H$ is closed under finite intersection. Therefore, one can construct a decreasing sequence

$$N_1 \supsetneq N_2 \supsetneq N_3 \supsetneq \cdots \supsetneq N_k \supsetneq \cdots$$

in $\mathfrak{N}_H$, hence each finite group G/N_i is a nontrivial homomorphic image of G/N_{i+1} ($i \in \omega$). Then let $K = \bigcap_{i \in \omega} N_i$, and using Lemma 15 for the group H with $\mathfrak{M} = \{N_i : i \in \omega\}$ we obtain

$$H/K \simeq \varprojlim_{i \in \omega} H/N_i.$$

We have a quotient H/K of H which is an inverse limit of an infinite sequence of finite groups, where the size of the groups is a strictly increasing sequence.

Let $S' \leq H/K$ be a null but non-meager subgroup (such an S' exists by Proposition 25, or 27). Let S be the preimage of S' under the canonical projection $\varphi_K : H \rightarrow H/K$

$$S = \varphi_K^{-1}(S') \leq H.$$

Using Corollary 23, S is null in H , but then since H is an open subgroup of G , S is null in G (the restriction of a Haar measure of G to an open subgroup H is a Haar measure of H , that is unique up to a positive multiplicative constant).

Then we can apply Lemma 20, i.e. S is not meager in the open subgroup H , thus is not meager in G . $\square$

3. MEAGER BUT NON-NULL SUBGROUPS

The following theorem due to Friedman can be found in [5]:

Theorem 30. *In the Cohen model (i.e. adding ω_2 Cohen reals to a model of $ZFC + CH$) the following holds:*

$$\begin{aligned} & \forall H \subseteq 2^\omega \times 2^\omega, \text{ if } H \text{ is } F_\sigma : \\ & (\exists C \times D \subseteq H, C \times D \notin \mathcal{N}) \Rightarrow (\exists A \times B \subseteq H \text{ measurable} : \mu(A \times B) > 0) \end{aligned}$$

Remark 31. *If X is a Polish space, and μ is a σ -finite Borel measure, then for any rectangle $C \times D \subseteq X \times X$, it is non-null (wrt. the product measure $\mu \times \mu$) iff $C \notin \mathcal{N}_\mu$ and $D \notin \mathcal{N}_\mu$.*

Next we prove that this property implies that every meager subgroup of a locally compact Polish group is of measure zero, this lemma is also from [5] (stating it only for $G = 2^\omega$, but the proof is the same).

Lemma 32. *Let G be a locally compact Polish group. Assume that the following holds:*

$$\begin{aligned} \forall H \subseteq G \times G, \text{ if } H \text{ is } F_\sigma : \\ (\exists C \times D \subseteq H, C \times D \notin \mathcal{N}) \Rightarrow (\exists A \times B \subseteq H \text{ measurable} : \mu(A \times B) > 0) \end{aligned} \quad (55)$$

Then every meager subgroup of G is null.

Proof. Let $S \leq G$ be a meager subgroup, assume on the contrary that S is non-null. Let $(H_i)_{i \in \omega}$ be nowhere dense closed subsets such that $\bigcup_{i \in \omega} H_i \supseteq S$. Now, if $m : G \times G \rightarrow G$ denotes the multiplication function, then $m^{-1}(\bigcup_{i \in \omega} H_i)$ is an F_σ set, containing $S \times S$, which is non-null. But then there is a measurable rectangle $A \times B \subseteq m^{-1}(\bigcup_{i \in \omega} H_i)$, which is of positive measure, thus by our Remark 31, $\mu(A), \mu(B) > 0$. Then, due to a Steinhaus type theorem [4], $AB = m(A \times B)$ has nonempty interior. But $AB \subseteq \bigcup_{i \in \omega} H_i$ which is meager, a contradiction (by the Baire category theorem). $\square$

First (in Lemma 34) we show that if every F_σ set $H \subseteq 2^\omega \times 2^\omega$ containing a rectangle of positive outer measure contains a measurable rectangle of positive measure, then this holds for arbitrary locally compact Polish groups. This yields that it is consistent with *ZFC* that in a locally compact Polish group meager subgroups are always null.

Later in Theorem 36 we reduce the general locally compact case to the case of Polish locally compact groups, by showing that if in locally compact Polish groups meager subgroups are null, then this is true in every locally compact group.

Remark 33. *If X is a Polish space, ν is a σ -finite Borel measure on X , then for every measurable set H there exists a Borel B such that $H \triangle B$ is null (wrt. μ).*

Lemma 34. *Assume that condition (55) holds in 2^ω (i.e. every F_σ subset of $2^\omega \times 2^\omega$ which contains a non-null rectangle must contain a measurable Haar-positive rectangle). Then (55) holds in every non-discrete locally compact Polish group G .*

Proof. Let μ denote the left Haar measure on G , and let ν denote the Haar measure on 2^ω . Since locally compact Polish groups are σ -compact, the Haar measure is σ -finite.

Claim 35. *There is a sequence of pairwise disjoint compact sets C_i ($i \in \omega$) in 2^ω , for which $\nu(2^\omega \setminus \bigcup_{i \in \omega} C_i) = 0$, and similarly a sequence of compact sets K_i ($i \in \omega$) in G with $\mu(K_i) > 0$ and $\mu(G \setminus \bigcup_{i \in \omega} K_i) = 0$, and there exist homeomorphisms $f_i : C_i \rightarrow K_i$, and positive constants r_i such that*

$$\forall B \subseteq C_i \text{ Borel} : \nu(B) = r_i \mu(f_i(B)).$$

Proof. First, using the inner regularity, and the σ -compactness of G , let E_i ($i \in \omega$) be a sequence of pairwise disjoint compact sets in G , for which $\mu(E_j) > 0$, and $\mu(G \setminus \bigcup_{i \in \omega} E_i) = 0$. Similarly, let F_i be a sequence of pairwise disjoint compact subsets of 2^ω such that $\nu(2^\omega \setminus \bigcup_{i \in \omega} F_i) = 0$, and $\nu(F_i) > 0$ ($\forall i$).

Now, since G is non-discrete, every open set is infinite in G , and an open set U with compact closure is thus an infinite set. But compact sets have finite measure. Then, by the invariance of the measure, every point has the same measure, that must be 0 because there exists infinite sets with finite measure, thus we obtain that μ is a continuous measure. We can apply the isomorphism theorem for measures [11, Thm 17.41.]. Let $g_i : F_i \rightarrow E_i$ be a Borel isomorphism (a bijection which is Borel, and so is its inverse), for which there exists $r_i > 0$ such that

$$\nu(H) = r_i \mu(f_i(H)) \quad (H \subseteq F_i \text{ Borel})$$

Now, by Lusin's theorem (see [11, Thm 17.12]) for every $H \subseteq C_i$ and $\varepsilon > 0$, the function $g_i|_H$ is continuous on a compact subset $H' \subseteq H$ with $\nu(H') > \nu(H) - \varepsilon$. Using this, there are pairwise disjoint compact subsets $(F_i^j)_{j \in \omega}$ in F_i , such that $g_i|_{F_i^j}$ is continuous, and $\nu(F_i \setminus \bigcup_{j \in \omega} F_i^j) = 0$. Let the sequence C_i ($i \in \omega$) be the enumeration of the F_i^j -s in type ω , and let $f_i = g_k^j$, if $C_i = F_k^j$, then choosing K_i to be $f_i(C_i)$ works. $\square$

Let $C = \bigcup_{i \in \omega} C_i \subseteq 2^\omega$, $K = \bigcup_{i \in \omega} K_i \subseteq G$, $f = \bigcup_{i \in \omega} f_i$, then

$$f : C \rightarrow K$$

is a bijection which is almost everywhere defined in 2^ω , and

- for each $S \subseteq C \subseteq 2^\omega$, if $\nu(S) = 0$, then $\mu(f(S)) = 0$,
- if S is measurable, $\nu(S) > 0$, then $f(S)$ is measurable, $\mu(f(S)) > 0$ (by our Remark 33, S differs from a Borel by a null set, f^{-1} is a Borel function, and f maps a null set to a null set),
- for every F_σ set $H \subseteq G$, $f^{-1}(H)$ is also F_σ (in 2^ω).

Now let $H \subseteq G \times G$ be an F_σ subset, and $D \times E \subseteq H$ be a rectangle of positive outer measure. Then $H \cap (K \times K)$ is still an F_σ set, and by Remark 31, $(D \times E) \cap (K \times K) = (D \cap K) \times (E \cap K)$ is of positive outer measure, since $\mu(G \setminus K) = 0$. Thus from now on, we can assume that $H \subseteq K \times K$: it suffices to find measurable non-null rectangles in such H -s. But $(f \times f)^{-1}(H)$ is F_σ , since $f \times f = (\bigcup_{i \in \omega} f_i) \times (\bigcup_{i \in \omega} f_i) = \bigcup_{i,j \in \omega} f_i \times f_j$ where the f_i -s are homeomorphisms between compact sets, so are the $f_i \times f_j$ -s. And $(f \times f)^{-1}(D \times E) = f^{-1}(D) \times f^{-1}(E)$ is a product of sets of positive outer measure, since f maps a nullset to a nullset. But then $(f \times f)^{-1}(H) \subseteq 2^\omega \times 2^\omega$, that contains $(f \times f)^{-1}(D \times E)$ contains a measurable $A \times B$ with $\nu(A), \nu(B) > 0$. Now, using that f maps a positive, measurable set to a positive, measurable set, we obtain that $\mu(f(A)), \mu(f(B)) > 0$, $f(A) \times f(B) \subseteq H$, hence H contains a non-null measurable rectangle, indeed. $\square$

Theorem 36. *If in every non-discrete locally compact Polish group meager subgroups are null, then this is true in non-discrete locally compact groups as well.*

Proof. Let $S \leq G$ be a meager but non-null subgroup. Then, applying Lemma 12, we get an open subgroup $H \leq G$, in which for each open neighborhood U of the identity, there is a compact normal subgroup $N_U \subseteq U$ such that G/N_U is a Lie group with finitely many connected components.

Then, since H is open, $S \cap H \leq H$ is clearly meager (in H).

Thus we can apply Lemma 19 for the co-meager set $R = H \setminus S$ in H and $J_0 = \emptyset$, yielding a compact normal subgroup $K' \triangleleft H$, and a co-meager set $R' \subseteq H/K'$, for which H/K' is a Polish space, and $\varphi_{K'}^{-1}(R') \subseteq R$, $(\varphi_{K'} : H \rightarrow H/K')$. Thus

$$\varphi_{K'}^{-1}(R') \cap (S \cap H) = \emptyset,$$

therefore

$$S \cap H \subseteq H \setminus \varphi_{K'}^{-1}(R') = \varphi_{K'}^{-1}((H/K') \setminus R'),$$

thus projecting $S \cap H$ to H/K' yields a subgroup in H/K' that is disjoint from the co-meager set R' .

We obtain that $\varphi_{K'}(S \cap H) \leq H/K'$ is a meager subgroup in a locally compact Polish group. Since in a discrete space a nowhere dense set must be empty, but we found a meager subgroup ($e_{H/K'} \in \varphi_{K'}(S \cap H)$), thus H/K' can not be discrete. Then $\varphi_{K'}(S \cap H) \leq H/K'$ is null by our assumptions (wrt. the Haar measure of H/K'). But then, by Corollary 23, the pull-back of $\varphi_{K'}(S \cap H)$ is also null in H

$$\mu_H(\varphi^{-1}(\varphi(S \cap H))) = 0,$$

and $S \cap H \subseteq \varphi^{-1}(\varphi(S \cap H))$, thus $\mu_H(S \cap H) = 0$.

Since H is an open subgroup of G , $S \cap H$ is null with respect to the Haar measure of G . But then let L be a set containing exactly one point from each left coset of H : $G = \bigcup_{l \in L} lH$ (i.e. it is a partition of G into pairwise disjoint open subsets). This partition gives rise to a partition of S : $S = \bigcup_{l \in L} ((lH) \cap S)$. Hence, by the left-invariance of μ_G , $(lH) \cap S$ is null, thus S is locally null. Now, from the fact that the Haar measure is inner regular wrt. compact sets, we can conclude that S is null in G .

□

Corollary 37. *It is consistent with ZFC that there are no meager subgroups of positive outer measure in locally compact groups.*

Example 38. *In the case of large, non- σ -compact locally compact groups, requiring outer regularity with respect to open sets, but the inner regularity with respect to compact sets holds only for open sets may result in a different measure than what we used. Consider the following example: take the reals with the discrete topology ($\mathbb{R}_d$), and with the euclidean topology ($\mathbb{R}$), then in $\mathbb{R}_d \times \mathbb{R}$:*

$S = \mathbb{R} \times \{0\}$ is a meager subgroup (it is nowhere dense, since there is a base consisting of sets of the form $\{r\} \times U$, $U \subseteq \mathbb{R}$ open). But any open U covering of

S can be partitioned into continuum many pairwise disjoint nonempty open sets: $U = \bigcup_{r \in \mathbb{R}} U \cap (\{r\} \times \mathbb{R})$, thus U is of infinite measure (recall that open sets have positive measure). Now, from the outer regularity wrt. open sets, S is of infinite measure.

Question 39. What can we say about small-large subgroups of not locally compact Polish groups, replacing "null wrt. the Haar measure" by "Haar-null" in the sense of Christensen? Do (always) exist meager but not Haar-null, and not Haar-null but meager subgroups in not locally compact Polish groups? (A subset X of a Polish group is Haar-null, if there is a Borel probability measure μ on G , and a Borel set $B \supseteq X$, such that for each $g, h \in G$ $\mu(gBh) = 0$ [6].)

REFERENCES

- [1] Arkhangel'skii, A., Tkachenko, M.: *Topological Groups and Related Structures* Atlantis Studies in Mathematics, Vol. 1, Atlantis Press, 2008.
- [2] Balka, R., Buczolich, Z. and Elekes, M.: *Topological Hausdorff dimension and level sets of generic continuous functions on fractals*. Chaos Solutions Fractals 45, pp 1579-1589., 2012.
- [3] Bartoszyński, T., Judah, H.: *Set Theory. On the Structure of the Real Line*. A K Peters, 1995.
- [4] Beck A., Corson H. H., Simon A. B. *The interior points of the product of two subsets of a locally compact group*. Proc. Amer. Math. Soc. 9, pp 648-652., 1958.
- [5] Burke, M. R.: *A theorem of Friedman on rectangle inclusion and its consequences*. Note of March 7, 1991.
- [6] Christensen, J.P.R.: *On sets of Haar measure zero in abelian Polish groups*. Israel J. Math. 13, pp 255-260., 1972.
- [7] Elekes, M., Tóth, Á.: *Covering locally compact groups by less than 2^ω many translates of a compact nullset*. Fund. Math. 193, pp 243-257., 2007.
- [8] Fremlin, D. H.: *Measure Theory: Volume 4, Topological Measure Spaces* 2003.
- [9] Gleason, A.M.: *The structure of locally compact groups*. Duke Math J. 18, pp 85-104., 1951.
- [10] Halmos, P. R.: *Measure theory*. Graduate Texts in Mathematics, 18, Springer-Verlag, 1974.
- [11] Kechris, A. S.: *Classical Descriptive Set Theory*. Graduate Texts in Mathematics, 156, Springer-Verlag, 1995.
- [12] Lee, J. M.: *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics, 218, Springer-Verlag, 2012.
- [13] Ribes, L., Zalesski, P.: *Profinite groups*. A Series of Modern Surveys in Mathematics, Vol. 40, Springer-Verlag, 2010.
- [14] Rosłanowski, A., Shelah, S.: *Small-large subgroups of the reals*. arXiv:1605.02261, 2016.
- [15] Tao, Terence: *Hilbert's Fifth Problem, and Related topics*. Graduate Studies in Mathematics, Vol. 153, American Mathematical Society, 2014.

EÖTVÖS LORÁND UNIVERSITY, INSTITUTE OF MATHEMATICS, PÁZMÁNY PÉTER S. 1/C, 1117 BUDAPEST, HUNGARY

E-mail address: sokmark@cs.elte.hu